

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
(International Bureau)

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: 7:

C12N 1/554, 1/555, 9/12, 9/16, C07K
16/18, 16/60, C12Q 1/68, A61K 38/45,
38/46

A2

(11) International Publication Number: WO 00/55333

(43) International Publication Date: 21 September 2000 (21/09/00)

(21) International Application Number: PCT/US99/0277

(22) International Filing Date: 17 March 2000 (17/03/00)

(23) Priority Date: 13 March 1999 (13/03/99)

(24) Priority Date: 13 March 1999 (13/03/99)

(25) Priority Date: 13 March 1999 (13/03/99)

(26) Priority Date: 13 March 1999 (13/03/99)

(27) Priority Date: 13 March 1999 (13/03/99)

(28) Priority Date: 13 March 1999 (13/03/99)

(29) Priority Date: 13 March 1999 (13/03/99)

(30) Priority Date: 13 March 1999 (13/03/99)

(31) Priority Date: 13 March 1999 (13/03/99)

(32) Priority Date: 13 March 1999 (13/03/99)

(33) Priority Date: 13 March 1999 (13/03/99)

(34) Priority Date: 13 March 1999 (13/03/99)

(35) Priority Date: 13 March 1999 (13/03/99)

(36) Priority Date: 13 March 1999 (13/03/99)

(37) Priority Date: 13 March 1999 (13/03/99)

(38) Priority Date: 13 March 1999 (13/03/99)

(39) Priority Date: 13 March 1999 (13/03/99)

(40) Priority Date: 13 March 1999 (13/03/99)

(41) Priority Date: 13 March 1999 (13/03/99)

(42) Priority Date: 13 March 1999 (13/03/99)

(43) Priority Date: 13 March 1999 (13/03/99)

(44) Priority Date: 13 March 1999 (13/03/99)

(45) Priority Date: 13 March 1999 (13/03/99)

(46) Priority Date: 13 March 1999 (13/03/99)

(47) Priority Date: 13 March 1999 (13/03/99)

Published
upon receipt of first report.

(54) TITLE: REGULATION OF INTRACELLULAR PHOSPHORYLATION

(57) Abstract

The invention provides human regulators of intracellular phosphorylation (IRIP) and polypeptides which identify and modulate IRIP. The invention also provides expression vectors, host cells, antibodies, agents, and analogs. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of IRIP.

(71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. (US005:3160) Power Drive, Palo Alto, CA 94304 (US).

(72) Inventor(s): 69135-068 (CN) 20 May 2000 (05/07/99) 9 July 1999 (09/07/99) 18 March 1999 (18/03/99)

(73) Agent(s): 11461-ET-CO., Dava et al., Incept Pharmaceuticals, Inc., 3160 Power Drive, Palo Alto, CA 94304 (US).

(81) Designated States: AE, AF, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CL, CN, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, MY, NZ, PE, PG, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, SM, SN, SR, ST, SV, SZ, TD, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VE, VU, WO, YU, ZA, ZW. ARIPO patent (CH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW). Eurasian patent (AM, AZ, BY, KG, KZ, KP, KZ, LU, RU, UA). OAPI patent (BF, BI, CF, CI, CM, GN, GW, ML, MR, NE, NG, TD, TO).

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the four pages of pamphlet publishing international applications under the PCT.

AE	Armenia	AF	Afghanistan	AL	Albania	AM	Armenia	AN	Andorra	AO	Angola	AR	Argentina	AT	Austria	AU	Australia	AX	Åland Islands	BA	Bosnia and Herzegovina	BB	Barbados	BC	Bahamas	BD	Bangladesh	BE	Belgium	BF	Burkina Faso	BG	Bulgaria	BH	Bahrain	BI	Burundi	BJ	Benin	BK	Belize	BL	Bolivia	BM	Bermuda	BN	Brunei Darussalam	BO	Bolivia	BR	Brazil	BS	Bahamas	BT	Bhutan	BV	Bouvet Island	BW	Botswana	BY	Belarus	BZ	Belize	CA	Canada	CC	Cocos (Keeling) Islands	CD	Congo	CE	Czechia	CF	Congo	CG	Congo	CH	Switzerland	CI	Côte d'Ivoire	CJ	Croatia	CK	Christmas Island	CL	Chile	CM	Camerun	CN	China	CO	Colombia	CR	Costa Rica	CS	Serbia and Montenegro	CU	Cuba	CY	Cyprus	CZ	Czechia	DD	East Germany	DE	Germany	DF	French Polynesia	DG	Dominican Republic	DH	Algeria	DI	Dominica	DJ	Dominican Republic	DK	Denmark	DM	Dominica	DO	Dominican Republic	DR	Dominican Republic	DU	Dubai	DZ	Algeria	EA	Spain	EC	Ecuador	EE	Estonia	EG	Egypt	EH	Western Sahara	ES	Spain	ET	Ethiopia	EU	European Union	FI	Finland	FJ	Fiji	FK	Falkland Islands	FM	Micronesia	FO	Faroe Islands	FR	France	GA	Gabon	GB	United Kingdom	GD	Grenada	GE	Georgia	GF	French Guiana	GG	Guernsey	GH	Ghana	GI	Gibraltar	GL	Greenland	GM	Gambia	GN	Guinea	GP	Guadeloupe	GQ	Equatorial Guinea	GR	Greece	GS	South Georgia and the South Sandwich Islands	GT	Guatemala	GU	Guam	GW	Guinea-Bissau	GY	Guyana	HA	Haiti	HC	Hong Kong	HD	North Korea	HE	Holaland	HF	French Polynesia	HG	Hong Kong	HH	Hong Kong	HI	Hawaii	HK	Hong Kong	HL	Hong Kong	HM	Hong Kong	HN	Honduras	HO	Hong Kong	HR	Croatia	HT	Haiti	HU	Hungary	IA	Iran	IB	Iran	IC	Iceland	ID	Indonesia	IE	Ireland	IL	Israel	IM	Isle of Man	IN	India	IO	Indian Ocean Territory	IQ	Iraq	IR	Iran	IS	Iceland	IT	Italy	JE	Jersey	JK	Japan	JM	Jamaica	JO	Jordan	JP	Japan	KA	Kazakhstan	KB	Kazakhstan	KC	Kazakhstan	KE	Kenya	KF	Kazakhstan	KG	Kyrgyzstan	KH	Kampuchea	KI	Kiribati	KJ	Kazakhstan	KK	Kazakhstan	KL	Kazakhstan	KM	Comoros	KN	St. Kitts and Nevis	KO	Korea	KP	North Korea	KR	South Korea	KS	Kazakhstan	KU	Kuwait	KV	Kazakhstan	KW	Kuwait	KX	Kazakhstan	KY	Cayman Islands	KZ	Kazakhstan	LA	Laos	LB	Lebanon	LC	St. Lucia	LD	Laos	LE	Lebanon	LF	Laos	LG	Laos	LH	Laos	LI	Liechtenstein	LJ	Laos	LK	Sri Lanka	LL	Laos	LM	Laos	LN	Laos	LO	Laos	LP	Laos	LR	Laos	LS	Lesotho	LT	Lithuania	LU	Luxembourg	LV	Latvia	LY	Libya	MA	Morocco	MB	Morocco	MC	Monaco	MD	Moldova	ME	Montenegro	MF	French Polynesia	MG	Madagascar	MH	Marshall Islands	MI	Maldives	MJ	Maldives	MK	Macedonia	ML	Mali	MM	Myanmar	MN	Mongolia	MO	Macao	MP	Marshall Islands	MQ	Martinique	MR	Morocco	MS	Montserrat	MT	Malta	MU	Mauritius	MV	Maldives	MW	Malawi	MX	Mexico	MY	Malaysia	MZ	Mozambique	NA	Namibia	NC	New Caledonia	NE	Niger	NF	Norfolk Island	NG	Nigeria	NH	Norway	NI	Nicaragua	NJ	Norway	NK	Norway	NL	Netherlands	NO	Norway	NP	Nepal	NR	Nauru	NT	Norway	NU	Niue	NV	Norway	NW	Norway	NY	Norway	NZ	New Zealand	OA	Oman	OB	Oman	OC	Oman	OD	Oman	OE	Oman	OF	Oman	OG	Oman	OH	Oman	OI	Oman	OJ	Oman	OK	Oman	OL	Oman	OM	Oman	ON	Oman	OO	Oman	OP	Oman	OQ	Oman	OR	Oman	OS	Oman	OT	Oman	OU	Oman	OV	Oman	OW	Oman	OX	Oman	OY	Oman	OZ	Oman	PA	Panama	PB	Panama	PC	Panama	PD	Panama	PE	Peru	PF	Panama	PG	Papua New Guinea	PH	Philippines	PI	Panama	PJ	Panama	PK	Pakistan	PL	Poland	PM	Puerto Rico	PN	Panama	PQ	Panama	PR	Puerto Rico	PS	Palestine	PT	Portugal	PV	Puerto Rico	PW	Panama	PX	Panama	PY	Paraguay	PZ	Panama	QA	Qatar	QB	Qatar	QC	Qatar	QD	Qatar	QE	Qatar	QF	Qatar	QG	Qatar	QH	Qatar	QI	Qatar	QJ	Qatar	QK	Qatar	QL	Qatar	QM	Qatar	QN	Qatar	QO	Qatar	QP	Qatar	QR	Qatar	QS	Qatar	QT	Qatar	QU	Qatar	QV	Qatar	QW	Qatar	QX	Qatar	QY	Qatar	QZ	Qatar	RA	Romania	RB	Romania	RC	Romania	RD	Romania	RE	Reunion	RF	Romania	RG	Romania	RH	Romania	RI	Romania	RJ	Romania	RK	Romania	RL	Romania	RM	Romania	RN	Romania	RO	Romania	RP	Romania	RS	Romania	RT	Romania	RU	Russia	RV	Romania	RW	Rwanda	RY	Romania	RZ	Romania	SA	Saudi Arabia	SB	Saint Helena	SC	Seychelles	SD	Sudan	SE	Sweden	SF	Sweden	SG	Singapore	SH	St. Helena	SI	Slovenia	SJ	Svalbard and Jan Mayen	SK	Slovakia	SL	Sierra Leone	SM	San Marino	SN	Senegal	SO	Somalia	SP	Spain	SQ	Slovenia	SR	Sri Lanka	SS	Sudan	ST	St. Vincent and the Grenadines	SV	El Salvador	SW	Swaziland	SX	Sri Lanka	SY	Syria	SZ	Swaziland	TA	Taiwan	TB	Taiwan	TC	Turks and Caicos Islands	TD	Togo	TE	Taiwan	TF	Taiwan	TG	Togo	TH	Thailand	TI	Taiwan	TJ	Tajikistan	TK	Tonga	TL	Timor-Leste	TM	Turkmenistan	TN	Tunisia	TO	Tonga	TR	Turkey	TS	Taiwan	TT	Trinidad and Tobago	TU	Turkmenistan	TV	Tuvalu	TW	Taiwan	TX	Taiwan	TY	Taiwan	TZ	Tanzania	UA	Ukraine	UB	Ukraine	UC	Ukraine	UD	Ukraine	UE	Ukraine	UF	Ukraine	UG	Uganda	UH	Ukraine	UI	Ukraine	UJ	Ukraine	UK	United Kingdom	UL	Ukraine	UM	Ukraine	UN	Ukraine	UO	Ukraine	UR	Ukraine	US	United States	UT	Ukraine	UU	Ukraine	UV	Ukraine	UW	Ukraine	UX	Ukraine	UY	Uruguay	UZ	Uzbekistan	VA	Vatican City	VB	Vatican City	VC	St. Vincent and the Grenadines	VE	Venezuela	VF	Venezuela	VG	Virgin Islands	VH	Venezuela	VI	Virgin Islands	VJ	Venezuela	VK	Venezuela	VL	Venezuela	VM	Venezuela	VN	Vietnam	VO	Vietnam	VP	Vietnam	VQ	Vietnam	VR	Vietnam	VS	Vietnam	VT	Vietnam	VU	Vanuatu	VX	Vietnam	VY	Vietnam	VZ	Vietnam	WA	Wallis and Futuna	WB	Wallis and Futuna	WC	Wallis and Futuna	WD	Wallis and Futuna	WE	Wallis and Futuna	WF	Wallis and Futuna	WG	Wallis and Futuna	WH	Wallis and Futuna	WI	Wallis and Futuna	WJ	Wallis and Futuna	WK	Wallis and Futuna	WL	Wallis and Futuna	WM	Wallis and Futuna	WN	Wallis and Futuna	WO	Wallis and Futuna	WP	Wallis and Futuna	WQ	Wallis and Futuna	WR	Wallis and Futuna	WS	Wallis and Futuna	WT	Wallis and Futuna	WU	Wallis and Futuna	WV	Wallis and Futuna	WW	Wallis and Futuna	WX	Wallis and Futuna	WY	Wallis and Futuna	WZ	Wallis and Futuna	XA	Xenia	XB	Xenia	XC	Xenia	XD	Xenia	XE	Xenia	XF	Xenia	XF	Xenia	XG	Xenia	XH	Xenia	XI	Xenia	XJ	Xenia	XK	Xenia	XL	Xenia	XM	Xenia	XN	Xenia	XO	Xenia	XP	Xenia	XQ	Xenia	XR	Xenia	XS	Xenia	XT	Xenia	XU	Xenia	XV	Xenia	XW	Xenia	XY	Xenia	XZ	Xenia	YA	Yemen	YB	Yemen	YC	Yemen	YD	Yemen	YE	Yemen	YF	Yemen	YG	Yemen	YH	Yemen	YI	Yemen	YJ	Yemen	YK	Yemen	YL	Yemen	YM	Yemen	YN	Yemen	YO	Yemen	YP	Yemen	YQ	Yemen	YR	Yemen	YS	Yemen	YT	Yemen	YU	Yugoslavia	YV	Yugoslavia	YW	Yugoslavia	YZ	Yugoslavia	ZA	South Africa	ZB	South Africa	ZC	South Africa	ZD	South Africa	ZE	South Africa	ZF	South Africa	ZG	South Africa	ZH	South Africa	ZI	South Africa	ZJ	South Africa	ZK	South Africa	ZL	South Africa	ZM	South Africa	ZN	South Africa	ZO	South Africa	ZP	South Africa	ZQ	South Africa	ZR	South Africa	ZS	South Africa	ZT	South Africa	ZU	South Africa	ZV	South Africa	ZW	South Africa	ZX	South Africa	ZY	South Africa	ZZ	South Africa	AA	Antarctica	AB	Antarctica	AC	Antarctica	AD	Antarctica	AE	Antarctica	AF	Antarctica	AG	Antarctica	AH	Antarctica	AI	Antarctica	AJ	Antarctica	AK	Antarctica	AL	Antarctica	AM	Antarctica	AN	Antarctica	AO	Antarctica	AP	Antarctica	AQ	Antarctica	AR	Antarctica	AS	Antarctica	AT	Antarctica	AU	Antarctica	AV	Antarctica	AW	Antarctica	AX	Antarctica	AY	Antarctica	AZ	Antarctica
----	---------	----	-------------	----	---------	----	---------	----	---------	----	--------	----	-----------	----	---------	----	-----------	----	---------------	----	------------------------	----	----------	----	---------	----	------------	----	---------	----	--------------	----	----------	----	---------	----	---------	----	-------	----	--------	----	---------	----	---------	----	-------------------	----	---------	----	--------	----	---------	----	--------	----	---------------	----	----------	----	---------	----	--------	----	--------	----	-------------------------	----	-------	----	---------	----	-------	----	-------	----	-------------	----	---------------	----	---------	----	------------------	----	-------	----	---------	----	-------	----	----------	----	------------	----	-----------------------	----	------	----	--------	----	---------	----	--------------	----	---------	----	------------------	----	--------------------	----	---------	----	----------	----	--------------------	----	---------	----	----------	----	--------------------	----	--------------------	----	-------	----	---------	----	-------	----	---------	----	---------	----	-------	----	----------------	----	-------	----	----------	----	----------------	----	---------	----	------	----	------------------	----	------------	----	---------------	----	--------	----	-------	----	----------------	----	---------	----	---------	----	---------------	----	----------	----	-------	----	-----------	----	-----------	----	--------	----	--------	----	------------	----	-------------------	----	--------	----	--	----	-----------	----	------	----	---------------	----	--------	----	-------	----	-----------	----	-------------	----	----------	----	------------------	----	-----------	----	-----------	----	--------	----	-----------	----	-----------	----	-----------	----	----------	----	-----------	----	---------	----	-------	----	---------	----	------	----	------	----	---------	----	-----------	----	---------	----	--------	----	-------------	----	-------	----	------------------------	----	------	----	------	----	---------	----	-------	----	--------	----	-------	----	---------	----	--------	----	-------	----	------------	----	------------	----	------------	----	-------	----	------------	----	------------	----	-----------	----	----------	----	------------	----	------------	----	------------	----	---------	----	---------------------	----	-------	----	-------------	----	-------------	----	------------	----	--------	----	------------	----	--------	----	------------	----	----------------	----	------------	----	------	----	---------	----	-----------	----	------	----	---------	----	------	----	------	----	------	----	---------------	----	------	----	-----------	----	------	----	------	----	------	----	------	----	------	----	------	----	---------	----	-----------	----	------------	----	--------	----	-------	----	---------	----	---------	----	--------	----	---------	----	------------	----	------------------	----	------------	----	------------------	----	----------	----	----------	----	-----------	----	------	----	---------	----	----------	----	-------	----	------------------	----	------------	----	---------	----	------------	----	-------	----	-----------	----	----------	----	--------	----	--------	----	----------	----	------------	----	---------	----	---------------	----	-------	----	----------------	----	---------	----	--------	----	-----------	----	--------	----	--------	----	-------------	----	--------	----	-------	----	-------	----	--------	----	------	----	--------	----	--------	----	--------	----	-------------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	------	----	--------	----	--------	----	--------	----	--------	----	------	----	--------	----	------------------	----	-------------	----	--------	----	--------	----	----------	----	--------	----	-------------	----	--------	----	--------	----	-------------	----	-----------	----	----------	----	-------------	----	--------	----	--------	----	----------	----	--------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	--------	----	---------	----	--------	----	---------	----	---------	----	--------------	----	--------------	----	------------	----	-------	----	--------	----	--------	----	-----------	----	------------	----	----------	----	------------------------	----	----------	----	--------------	----	------------	----	---------	----	---------	----	-------	----	----------	----	-----------	----	-------	----	--------------------------------	----	-------------	----	-----------	----	-----------	----	-------	----	-----------	----	--------	----	--------	----	--------------------------	----	------	----	--------	----	--------	----	------	----	----------	----	--------	----	------------	----	-------	----	-------------	----	--------------	----	---------	----	-------	----	--------	----	--------	----	---------------------	----	--------------	----	--------	----	--------	----	--------	----	--------	----	----------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	--------	----	---------	----	---------	----	---------	----	----------------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	------------	----	--------------	----	--------------	----	--------------------------------	----	-----------	----	-----------	----	----------------	----	-----------	----	----------------	----	-----------	----	-----------	----	-----------	----	-----------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	---------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------------------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	-------	----	------------	----	------------	----	------------	----	------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	--------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------	----	------------

REGULATORS OF INTRACELLULAR PHOSPHORYLATION

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of regulators of intracellular phosphorylation and to the use of these sequences in the diagnosis, treatment, and prevention of neurological, cell proliferative, and autoimmune/inflammatory disorders.

BACKGROUND OF THE INVENTION

Reversible protein phosphorylation is the main strategy for controlling the activities of eukaryotic cells. Kinases and phosphatases regulate reversible phosphorylation reactions, and thus critical components of intracellular signal transduction pathways. Protein kinases transfer the high energy phosphate from adenosine triphosphate (ATP) to specific protein targets in response to extracellular signals (such as hormones, neurotransmitters, and growth and differentiation factors), cell cycle checkpoints (for example, signals associated with mitotic entry), and environmental or nutritional stresses. Protein phosphatases mediate kinase effect by removing phosphate groups from proteins.

It is estimated that more than 1000 of the 10,000 proteins active in a typical mammalian cell are phosphorylated. In general, protein activity is switched by phosphorylation, and this is analogous to turning on a molecular switch. When the switch is turned on, the appropriate protein kinase activates a metabolic enzyme, regulatory protein, receptor, cytoskeletal protein, ion channel or pump, or transcription factor. Protein activity is repressed by dephosphorylation when down-regulation of a signalling pathway is required. The coordinate activities of kinases and phosphatases regulate key processes such as cell proliferation, cell differentiation, cell-cell communication, and cell cycle progression. Uncontrolled signalling has been implicated in a variety of disease conditions including inflammation, cancer, arteriosclerosis, and atherosclerosis.

Protein kinases phosphorylate protein acceptor molecules on hydroxylated amino acids. These kinases comprise the largest known protein group, a superfamily of enzymes with widely varied functions and specificities. Protein kinases are usually named after substrates, regulatory molecules, or some aspect of a mutant phenotype. With regard to substrates, protein tyrosine kinases are roughly divided into two groups: those that phosphorylate tyrosine residues (protein tyrosine kinases (PTKs)), and those that phosphorylate serine or threonine residues (serine/threonine kinase (STKs)). A few protein kinases have dual specificity and phosphorylate serine, threonine, and tyrosine residues. Some STKs and PTKs possess structural characteristics of both families (Hardie, G. and S. Hanke (1995) *The Protein Kinase Pocket Book*, Vol. 1:7-20. Academic Press, San Diego, CA).

Almost all kinases contain a conserved 250-300 amino acid kinase domain that folds into a

two-lobed structure. The primary structure of the kinase domain is conserved and can be further subdivided into 11 subdomains. The smaller N-terminal lobe of the kinase domain, which contains subdomains I through IV, is primarily involved in the binding and orientation of the ATP (or GTP) donor molecule. The larger C-terminal lobe, which contains subdomains VI through XI, binds the protein substrate and carries out transfer of the gamma phosphate from ATP to the hydroxyl group of a serine, threonine, or tyrosine residue. Subdomain V spans the two lobes. Each of the 11 subdomains contains specific amino acid residues and motifs that are characteristic of the particular subdomain and are highly conserved (Hardie, G. and S. Hands (1995) *The Protein Kinase Atlas*, Book 6, Vol. 1:7-20. Academic Press, San Diego CA). In particular, two protein kinase signature sequences have been identified in the kinase domain, the first containing an active site lysine residue involved in ATP binding (subdomain II), and the second containing an aspartate residue important for catalytic activity (subdomain VI). Kinases may also be categorized into families by the different amino acid sequences (generally between 5 and 100 residues) located on either side of, or inserted into, the kinase domain. These additional amino acid sequences are involved in the regulation of kinase activities and interactions with other proteins.

PTKs may be classified as either transmembrane or non-transmembrane proteins. Transmembrane PTKs function as receptors for most growth factors. Growth factors bind to the receptor tyrosine kinase (RTK), causing the RTK to phosphorylate itself (autophosphorylation) and specific intracellular second messenger proteins. Growth factors that bind RTKs include epidermal growth factor, platelet-derived growth factor, fibroblast growth factor, hepatocyte growth factor, insulin and insulin-like growth factors. Nerve growth factor, vascular endothelial growth factor, and macrophage colony stimulating factor

Non-transmembrane PTKs form signalling complexes with the cytosolic domains of plasma membrane receptors. Receptors that signal through non-transmembrane PTKs include receptors for cytokines and hormones (e.g. growth hormone and prolactin), and antigen-specific receptors on T and B lymphocytes. Many non-transmembrane PTKs were first identified as the products of mutant oncogenes in cancer cells in which PTK activation was no longer subject to normal cellular controls. About one third of the known oncogenes encode PTKs, and it is well known that cellular transformation (oncogenesis) is often accompanied by increased tyrosine phosphorylation activity (Charbonneau H and N. K. Tonks (1992) *Annu. Rev. Cell Biol.* 8:463-499). Regulation of PTK activity may therefore be an important stepway in controlling some types of cancer.

STKs are non-transmembrane proteins. STKs include second messenger-dependent protein kinases, which primarily mediate the effects of second messengers such as cyclic AMP, cyclic GMP, inositol triphosphate, phosphatidylinositol 3,4,5-trisphosphate, cyclic ADP ribose, arachidonic acid, diacylglycerol, and calcium-calmodulin (CaM). STKs include cyclic AMP dependent protein kinases,

Protein phosphatases regulate the effects of protein kinases by removing phosphate groups from nucleotides activated by kinases. Phosphatases are characterized as either tyrosine-specific or serine/threonine-specific based on their preferred phospho-amino acid substrate, although some protein phosphatases have dual specificity for both serine/threonine and tyrosine groups.

5 Serine/threonine phosphatases dephosphorylate phosphoserine and phosphothreonine residues, and are important regulators of many cAMP-mediated hormone responses (Cohen, P. (1989) *Annu. Rev. Biochem.* 58:453-508). Serine/threonine phosphatases generally comprise two or more subunits and have broad and overlapping protein substrate specificities.

Tyrosine phosphatases are generally monomeric proteins which function primarily in the transduction of signals across the plasma membrane, and are categorized as either transmembrane receptor-like proteins or soluble non-transmembrane proteins. Tyrosine phosphatases reverse the effects of PTKs, removing phosphate groups from tyrosine residues of phosphorylated proteins, and play a significant role in cell cycle and cell signaling processes, lymphocyte activation, and cell adhesion (Charbonneau, H. and N.K. Tonks (1992) *Annu. Rev. Cell Biol.* 8:463-493). In the process of cell division, for example, a specific tyrosine phosphatase called M-phase inducer phosphatase plays a key role in the induction of mitosis by dephosphorylating and activating CDK2, a cell division-specific PTK (Nishida, S. et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:5139-5143).

Tyrosine phosphatases share a conserved catalytic domain of about 250 amino acids which contains the active site. The active site consensus sequence consists of 13 amino acids, including a cysteine residue that is essential for phosphatase activity. In addition, the gene encoding at least eight tyrosine phosphatases have been mapped to chromosomal regions that are translocated or rearranged in various neoplastic conditions, including lymphoma, leukemia, small cell lung carcinoma, adenocarcinoma, and neuroblastoma (Charbonneau, H. and N.K. Tonks (1992) *Annu. Rev. Cell Biol.* 8:463-493). As previously noted, many PTKs are encoded by oncogenes, and oncogenesis is often accompanied by increased tyrosine phosphorylation activity. It is therefore possible that tyrosine phosphatases may prevent or reverse cell transformation and the growth of various cancers by controlling the levels of tyrosine phosphorylation in cells. This hypothesis is supported by studies showing that overexpression of tyrosine phosphatases can suppress transformation in cells, and that specific inhibition of tyrosine phosphatases can enhance cell transformation (Charbonneau and Tonks, 1992).

In addition to protein phosphorylation, lipid phosphorylation also plays a role in certain signaling pathways. The phosphorylation of phosphatidylinositol is involved in activation of the PKC signaling pathway. Recently, a sphingolipid metabolite, sphingosine-1-phosphate (SPP), has emerged as a novel lipid second-messenger with both extracellular and intracellular actions (Kohama, T. et al. (1998) *J. Biol. Chem.* 273:23722-23728). Extracellularly, SPP is a ligand for the G-protein

coupled receptor EDG-1 (endothelial-derived, G-protein coupled receptor). Intracellularly, SPP regulates cell growth, survival, motility, and cytoskeletal changes. SPP levels are regulated by sphingosine kinases that specifically phosphorylate D-erythro-sphingosine to SPP. The importance of sphingosine kinase in cell signaling is indicated by the fact that various stimuli, including platelet-derived growth factor (PDGF), nerve growth factor, and activation of PKC, increase cellular levels of SPP by activation of sphingosine kinase, and the fact that competitive inhibitors of the enzyme selectively inhibit cell proliferation induced by PDGF (Kohama et al., 1993).

The discovery of new regulators of intracellular phosphorylation and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of neurological, cell proliferative, and autoimmune/inflammatory disorders.

SUMMARY OF THE INVENTION

The invention features purified polypeptides, regulators of intracellular phosphorylation, referred to collectively as "HRP" and individually as "HRP-1," "HRP-2," "HRP-3," "HRP-4," "HRP-5," "HRP-6," "HRP-7," "HRP-8," "HRP-9," "HRP-10," "HRP-11," "HRP-12," "HRP-13," and "HRP-14." In one aspect, the invention provides an isolated polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-14. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-14.

The invention further provides an isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-14. In one alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:1-14.

Alternatively, the polynucleotide is selected from the group consisting of SEQ ID NO:1-14. In one alternative, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, c) a biologically active fragment of an amino acid sequence selected

from the group consisting of SEQ ID NO:1-4, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-4. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO.1-14, b) a naturally occurring amino acid sequence having at least 95% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO.1-14, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO.1-14, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO.1-14. The method comprises a) an amino acid sequence selected from the group consisting of SEQ ID NO.1-14. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant plasmid encoding a precursor sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

15 Additionally, the invention provides a labeled antibody which specifically binds to a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO: 1, 4; b) a mutually occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 1-14; c) a nonlogically altered fragment of an amino acid sequence selected from the group consisting of SEQ ID NO: 1-14; d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO: 1-14.

The invention further provides an isolated polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:1-26, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:1-26, c) a polynucleotide sequence complementary to a), or d) a polynucleotide sequence complementary to b). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO.1-53; b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO.1-53; c) a polynucleotide sequence complementary to a) or d) a polynucleotide sequence complementary to b). The method comprises a) hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a polynucleotide sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed

between solid probe and solid target polynucleotide, and b) detecting the presence or absence of solid hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 30 contiguous nucleotides. In another alternative, the probe comprises at least 66 contiguous nucleotides.

- 5 The invention further provides a pharmaceutical composition comprising an effective amount of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, b) a mutually occurring amino acid sequence having in at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, and a pharmaceutically acceptable excipient. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional HARP.
- 10

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-14. (b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, c) a heterologous fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-14. The method comprises a) exposing a sample comprising the polypeptide to a

compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional HTRP, comprising administering to a patient in need of such treatment the pharmaceutical composition.

amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophobicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

5 The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, polypeptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and the terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

10 "Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technology, well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of HRP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of HRP either by directly interacting with HRP or by acting on components of the biological pathway in which HRP participates.

The term "antigen" refers to an immunoglobulin molecule as well as to fragments thereof, such as Fab, Fab₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind HRP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

20 The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antiserum" refers to any composition capable of base-pairing with the "sense" strand of a specific nucleic acid sequence. Antiserum compositions may include DNA, RNA, peptide nucleic acid (PNA), oligonucleotides having modified backbone linkages, such as phosphorothioates,

30 methyolphosphonates, or heavy phosphonates; oligonucleotides having modified sugar groups such as 2-methoxyethyl sugars or 2-methoxyvinyl sugars; or oligonucleotides having modified bases such as 5-substituted cytosine, 2-deoxyguanosine, or 7-deaza-2-deoxyguanosine. Antiserum molecules may be produced by any method including chemical synthesis or transfection. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic HRP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5'-A-G-T-3'" bonds to the complementary sequence "3'-T-C-A-5'". Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution.

25 Compositions comprising polynucleotide sequences encoding HRP or fragments of HRP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate, SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

A "consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncloned bases, extended using the XL-PCR kit (Pharman-Lerner, New York, CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of one or more large clones and, in some cases, one or more public domain ESTs, using a computer program for fragment assembly, such as the GEL VIEW fragment assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

35

methyolphosphonates, or heavy phosphonates; oligonucleotides having modified sugar groups such as 2-methoxyethyl sugars or 2-methoxyvinyl sugars; or oligonucleotides having modified bases such as 5-substituted cytosine, 2-deoxyguanosine, or 7-deaza-2-deoxyguanosine. Antiserum molecules may be produced by any method including chemical synthesis or transfection. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic HRP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5'-A-G-T-3'" bonds to the complementary sequence "3'-T-C-A-5'". Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

20 A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution.

25 Compositions comprising polynucleotide sequences encoding HRP or fragments of HRP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate, SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

A "consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncloned bases, extended using the XL-PCR kit (Pharman-Lerner, New York, CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of one or more large clones and, in some cases, one or more public domain ESTs, using a computer program for fragment assembly, such as the GEL VIEW fragment assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

30

35

WO 00/55333

PCT/US00/0777

(i.e., less than about 30% similarity or identity). In the absence of non-specific binding, the substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 1.1.2a sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1993) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weight" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequence pairs.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gov/blast2.html>.

The "BLAST 2 Sequences" tool can be used for both bias and bias (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool (Version 2.0.9 (May-97, 1999)) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62
Reward for match: 1
Penalty for mismatch: -3
Open Gap: 5 and Extension Gap: 2 penalties
Gap x drop-off: 50

15

WO 00/55333

PCT/US00/0777

Expect: 10
Word Size: 11
Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number; or may be measured over a shorter length, for example, over the length of a fragment chosen from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the hydrophobicity and activity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 1.1.2a sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool (Version 2.0.9 (May-97, 1999)) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62
Open Gap: 11 and Extension Gap: 1 penalties
Gap x drop-off: 50
Expect: 10
Word Size: 3

16

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for

5 instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to determine a length over which percentage identity may be measured.

10 "Human artificial chromosome" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term "humanized antibody" refers to antibody molecules in which the amino acid sequences in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

15 "Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of identity.

Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the

20 stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Generally, such wash temperatures are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook et al. 1989, *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

35 High stringency conditions for hybridization between polynucleotides of the present

invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%.

Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular

5 circumstances, such as for RNA-DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

10 The term "hybridization complex" refers to a complex, formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C₀ or R₀ analysis) or formed between one

15 nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide

sequence resulting in the addition of one or more amino acid residues, or nucleotides, respectively. "Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

20 An "immunogenic fragment" is a polypeptide or oligopeptide fragment of HRP which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of HRP which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

25 The terms "element" and "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

30 The term "module" refers to a change in the activity of HRP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of HRP.

35 The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or

synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide, nucleic acid (RNA), or to any DNA-labile or RNA-labile material.

- 10 "Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences may be in close proximity or contiguous and, where necessary, to join two protein coding regions, in the same reading frame.

- 15 "Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-sense agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcription elongation, and may be employed to extend their lifespan in the cell.

- 20 "Probe" refers to nucleic acid sequences encoding TRAP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule.

- Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

- 25 "Primer" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

- 30 Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

- 35 Methods for preparing and using probes and primers are described in the references, for example Sambrook et al., 1989, *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel et al., 1987, *Current Protocols in Molecular Biology*; Greenlee, Fohl, Arose, & Wiley-Interscience, New York NY; Imlis et al., 1990, *PCR: Practical A Guide to Methods and Applications*, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA). Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.16 software is useful for the selection of PCR primer pairs of up to

- 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kbases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the Primer3 primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas, TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "navigating library," in which

- 15 sequences to avoid a primer binding site are user-specified. Primer3 is, in particular, for the selection of oligonucleotide for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The Primer3 program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved

- 20 regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotide and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

- 25 A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, *et al.*, 1989. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

- 30 Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to transform a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

- 35 An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose

instead of deoxythymine.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acid encoding HRP1 or fragments thereof, or HRP1 itself, may comprise a bodily fluid; an extract from a cell; chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print, etc.

The terms "specific binding" and "specifically binding" refer to that intersection between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

"Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term

"transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor

of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or *in situ* zygote fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, and plants and animals. The isolated DNA of the present invention can be

introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989).

DEFINITIONS

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blast with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or

greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "spliced," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of

polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The

presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blast with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human regulators of intracellular phosphorylation (HRP1), the polynucleotides encoding HRP1, and the use of these compositions for

the diagnosis, treatment, or prevention of neurological cell proliferative, and autoimmune/inflammatory disorders.

Table 1 lists the incise clones used to assemble full length nucleotide sequences encoding HRP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the incise clones in which nucleic acids encoding each HRP were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows incise clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. The incise clones in column 5 were used to assemble the consensus nucleotide sequence of each HRP, and are useful as fragments in hybridization techniques.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologue sequences as identified by BLAST analysis; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding HRP. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification techniques to identify SEQ ID NOs 15-28 and to distinguish between SEQ ID NOs 15-28 and related polynucleotide sequences. For SEQ ID NOs 15-27, the polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express HRP as a fraction of total tissues expressing HRP. Column 4 lists diseases, disorders, or conditions associated with these tissues expressing HRP as a fraction of total tissues expressing HRP. Column 5 lists the vectors used to subclone each cDNA library.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding HRP were isolated. Column 1 references the nucleotide SEQ ID NOs; column 2 shows the cDNA libraries from which these clones were isolated; and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2. The invention also encompasses HRP variants. A preferred HRP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the HRP amino acid sequence, and which contains at least one functional or structural

WFO 00, 2
characteristic of HRP.

The invention also encompasses polynucleotides which encode HRP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NOs 15-28, which encodes HRP. The polynucleotide sequences of SEQ ID NOs 15-28, as presented in the Sequence Listing, enhance the equivalent RNA sequences, wherein occurrences of the antigenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding HRP. In particular, such a variant polynucleotide sequence will have at least about 80%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding HRP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NOs 15-28 which has at least about 80%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NOs 15-28. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of HRP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding HRP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention encompasses each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring HRP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode HRP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring HRP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding HRP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding HRP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode HRP and

synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding HRP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:15-28 and fragments thereof under various conditions of stringency. (See, e.g., Walli, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:307-311.) Hybridization conditions, including annealing and wash conditions, are described in "Hybridizations".

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), the thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequencing preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI/CATALYST 800 thermal cycler (Applied Biosystems, Foster City CA). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Perkin-Elmer), the MEGACASE 1000 DNA sequencing system (Molecular Dynamics, Warrington, UK), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) *Short Protocols in Molecular Biology*, John Wiley & Sons, New York, NY, pp. 717-765; Meyers, R.A. (1995) *Molecular Biology and Biochemistry*, Wiley & Sons, New York, NY, pp. 856-873.)

The nucleic acid sequences encoding HRIP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as primers and regulatory elements. For example, one method which may be employed

restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar G. (1993) PCR Methods, Appl. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8183.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Appl. 1:1-11.) In this method, multiple restriction enzyme

155/000

PCT/US00/07277

digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods would be used to retrieve unknown sequences are known in the art. (See, e.g., Pelzer, J.D. et al. (1997) *Nucleic Acids Res.* 25:3555-3560). Additionally, one may use PCR, nested primers, and PROMOTER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding insertion junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis Program (National Biosciences, Plymouth MN) or another appropriate program. About 21 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electronic signal using appropriate software (e.g., GENOTYPE and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amount in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode HRP may be cloned in recombinant DNA molecules that direct expression of HRP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express HRP.

The nucleotide sequence of the present invention can be engineered using methods generally known in the art in order to alter HRP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing and/or expression of the gene product. DNA shuffling by random fragmentation and PCR assembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction

sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

- The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MODULAR RECOMBINING (Maxygen Inc., Santa Clara CA, described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) *Nat. Biotechnol.* 17:793-797; Christians, F.C. et al. (1999) *Nat. Biotechnol.* 17:259-264; and Garmen, A. et al. (1996) *Nat. Biotechnol.* 14:315-319) to alter or improve the biological properties of HRP, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then resubstituted until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

- In another embodiment, sequences encoding HRP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Cantabers, M.H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-233.)

- Alternatively, HRP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Rodighiero, J.Y. et al. (1993) *Science* 260:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Foster/Eli Lilly). Additionally, the amino acid sequence of HRP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

- The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Rejzler (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptide may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) *Proteins, Structures and Molecular Properties*, W.H. Freeman, New York NY.)

- In order to express a biologically active HRP, the nucleotide sequence encoding HRP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and

- inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding HRP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding HRP. Such signals include the ATG initiation codon and adjacent sequences, e.g., the Kozak sequence. In cases where sequences encoding HRP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequences, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Schmitt, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

- Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding HRP and appropriate transcriptional and translational control elements. These methods include *in situ* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. (See, e.g., Sambrook, J. et al. (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

- A variety of expression vector/host systems may be utilized to contain and express sequences encoding HRP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., T7 or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

- In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding HRP. For example, routine cloning, subcloning, and integration of polynucleotide sequences encoding HRP can be achieved using a multifunctional E. coli vector such as pBLUESCRIPT (Stratagene, La Jolla CA) or pSPORT1 (pilot) (Life Technologies). Ligation of sequences encoding HRP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription-translation sequencing, single strand rescue with helper phage, and creation of needed deletions in the cloned sequence. (See, e.g., Van Hecke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5305-5309.) When large quantities of HRP are needed, e.g., for the production of

fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies relative to two non-interfering epitopes on HRP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) *Serological Methods, A Laboratory Manual*, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) *Current Protocols in Immunology*, Greene Pub. Associates and Wiley-Interscience, New York NY; and Ponder, J.D. (1998) *Immunotechnical Protocols*, Humana Press, Towon NJ)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled

- 10 hybridization or PCR probes for detecting sequences related to polynucleotides encoding HRP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequence encoding HRP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for
- 15 cases of detection include radionuclides, enzymes, fluorescent, chemoluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

- 20 Host cells transfected with nucleotide sequence encoding HRP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transfected cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode HRP may be designed to contain signal sequences which direct secretion of HRP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, phosphorylation,

- 30 phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "pro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas, VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

- 35 In another embodiment of the invention, animal, modified, or recombinant nucleic acid

- sequences encoding HRP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric HRP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of HRP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thiodiolin (Tdx), calmodulin binding peptide (CBP), c-Ha, FLAG, c-myc, and hemagglutinin (HA). GST, MBP, Tdx, CBP, and c-Ha enable purification of heterologous fusion proteins on immobilized glutathione, maltose, p-phenylene oxide, calmodulin, and metal-chelate resins, respectively. FLAG, c-myc, and hemagglutinin (HA) enable immunofluorescent purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the HRP encoding sequence and the heterologous protein sequence, so that HRP may be cleaved away from the heterologous moiety following purification.
- 15 Methods for fusion protein expression and purification are discussed in Ausubel (1995), supra, ch. 10. A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

- In a further embodiment of the invention, synthesis of radiolabeled HRP may be achieved in vitro using the T7T3 rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

- 25 Fragments of HRP may be produced not only by recombinant means, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, *SMBL*, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the ABI 431A peptide synthesizer (Pierce-Elmer). Various fragments of HRP may be synthesized separately and then combined to produce the full length molecule.

THEORETICALS

- Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of HRP and regulators of intracellular phosphorylation. In addition, the expression of HRP is closely associated with neurological tissue, with cancer and other cell proliferative disorders, and with inflammation and the immune response. Therefore, HRP appears to play a role in neurological, cell proliferative, and autoimmune/inflammatory disorders. In the treatment of disorders associated with increased HRP expression or activity, it is desirable to decrease the expression or activity of HRP. In the treatment of disorders associated with decreased HRP

expressive or activity, it is desirable to increase the expression or activity of HRP.

Therefore, in one embodiment, HRP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of HRP.

Examples of such disorders include, but are not limited to, a neurological disorder such as epilepsy.

- 5 is ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, atrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, solidary empyema, epidural abscess, suppurative intercranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease: prion diseases including bun, Creutzfeldt-Jakob disease, and Gerstmann-Sträussler-Scheinker syndrome; focal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebellofrenic hemangioblastomatosis, encephalotrigeminal syndrome, neural retardation and other developmental disorders of the central nervous system, cerebral palsy,
- 15 neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dysrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis; inherited, metabolic, endocrine, and toxic myopathies; myasthenia gravis, periodic paralysis; mental disorders including mood, anxiety, and schizophrenia disorders; seasonal affective disorder (SAD); anorexia, amnesia, cataplexy, diabetic neuropathy, tardive dyskinesia, dysomnia, paranoid psychosis, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia, a cell proliferative disorder such as actinic keratosis, actinocarcinoma, osteosarcoma, bursitis, arthritis, hepatitis, mixed connective tissue diseases (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and a cancer including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, a cancer of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, prostate, salivary gland, rect, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, skin, spleen, testis, thymus, thyroid, and uterus; and an
- 25 autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, alopecia areata, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dysrophy (APECED), bronchiectasis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocyteopenia, erythroblastosis fetalis, erythema nodosum, atrophic
- 35 gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's

thyroiditis, hyporesponsibility, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, paracitosis, polyomyositis, psoriasis, Reiter's disease, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extrasporal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma.

In another embodiment, a vector capable of expressing HRP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of HRP including, but not limited to, those described above.

- 10 In a further embodiment, a pharmaceutical composition comprising a substantially purified HRP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of HRP including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of HRP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of HRP including, but not limited to, those listed above.

In a further embodiment, an antagonist of HRP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of HRP. Examples of such disorders include, but are not limited to, those neurological, cell proliferative, and autoimmune/inflammatory disorders described above. In one aspect, an antibody which specifically binds HRP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express HRP.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding HRP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of HRP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of HRP may be produced using methods which are generally known in the art. In particular, purified HRP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind HRP. Antibodies to HRP may also

disclosed in the literature. (See, e.g., Offhand, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:295-299.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunochemical assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such

immunoassay typically involve the measurement of complex formation between HRP and its substrate. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering HRP epitopes is generally used, but a competitive binding assay may also be employed (Pound, 1982).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for HRP. Affinity is expressed as an equilibrium constant, K , which is defined as the molar concentration of HRP-antibody complex

The K_d determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple HRP epitopes, represents the average affinity, or avidity, of the antibodies for a HRP. The K_d determined for a preparation of monoclonal antibodies, which are monospecific for a particular HRP epitope, represents the measure of affinity. High-affinity antibody preparations with K_d ranging from 10^6 to 10^{12} mol/L are referred to as immunoreagents in which the

HRP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_d ranging from about 10^6 to 10^8 M are preferred for use in immunofluorescence and similar applications, whereas high-affinity preparations are preferred for use in immunoassays. Antibodies which ultimately require dissociation of HRP, preferably in active form, from the antibody (Cmyr, D. (1988) *Antibodies, Volume 1: A Practical Approach*, IRL Press, Wobington, DC).

Laddell, J. E. and Oyer, A. (1991) *A Practical Guide to Monoclonal Antibodies*, John Wiley & Sons, New York, NY.

The liver and spleen of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a collection of antibody preparations consisting of about 12 serum-specific, antibody-

preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of HRP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity and

guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Carby, *SUB2*, and Coligan et al. *SUB3*.)

In another embodiment of the invention, the polynucleotides encoding HRP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding HRP may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding HRP. Thus, complementary molecules or fragments may be used to modulate HRP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding HRP.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding HRP. (See, e.g., Sambrook, *SUB2*; Ausubel, 1995, *SUB3*.)

Genes encoding HRP can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding HRP. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5' or regulatory regions of the gene encoding HRP. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may be employed. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.J. Carr, *Molecular and Immunologic Approaches*, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme

molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered haemichannel motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding HRP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding HRP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2'-O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, quosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytosine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use *in vitro*, *in vivo*, and *ex vivo*. For *ex vivo* therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) *Nu. Biotechnol.* 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of HRP, antibodies to HRP, and mimetics, agonists, antagonists, or inhibitors of HRP. The compositions

- 5 may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

- 10 The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intracerebral, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

- In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of *Remington's Pharmaceutical Sciences* (Mack Publishing, Easton PA).

- Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

- Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally after grinding) to obtain tablets or dragee cores. Suitable excipients can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and vegetable, and protein, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginate acid or salt thereof, such as sodium alginate.

- Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbonyl gel, polyethylene glycol, and/or titanium dioxide, liquid solutions, and suitable organic solvents or solvent mixtures. Dyes and pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

- Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starch, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

- Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextrose. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, (glycerides, or liposomes. Non-lipid polymeric amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

- For topical or nasal administration, preparations appropriate to the particular barrier to be permeated are used in the formulation. Such preparations are generally known in the art. The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

- The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in aqueous or other protic solvents than are the corresponding free base forms. In other cases, the preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 3% sucrose, and 2% to 7% mannitol, in a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

- After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of IHRP, such labeling would include amount, frequency, and method of administration.

- Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

- For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, or pigs.

An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example HRP or fragments thereof, antibodies of HRP, and agonists, antagonist or inhibitors of HRP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED₅₀ (the dose therapeutically effective in 50% of the population) or LD₅₀ (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effect is the therapeutic index, which can be expressed as the LD₅₀/ED₅₀ ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED₅₀ with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 µg to 100,000 µg, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind HRP may be used for the diagnosis of disorders characterized by expression of HRP, or in assays to monitor patients being treated with HRP or agonists, antagonists, or inhibitors of HRP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for HRP include methods which utilize the antibody and a label to detect HRP in human body fluids

or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring HRP, including ELISAs, RIA, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of HRP expression. Normal or standard values for HRP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to HRP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of HRP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding HRP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of HRP may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of HRP, and to monitor regulation of HRP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding HRP or closely related molecules may be used to identify nucleic acid sequences which encode HRP. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding HRP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the HRP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:15-28 or from genomic sequences including promoters, enhancers, and introns of the HRP gene.

Means for producing specific hybridization probes for DNAs encoding HRP include the cloning of polynucleotide sequences encoding HRP or HRP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radioisotopes such as ³²P or ³⁵S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

43

In a particular aspect, the nucleotide sequences encoding HRP² may be useful in assaying that

for the treatment of an individual patient.

with the presence of a disorder

months.

A

of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding HRP

- 5 may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced *in situ*. Oligomers will preferably contain a fragment of a polynucleotide encoding HRP, or a fragment of a polynucleotide complementary to the polynucleotide encoding HRP, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or
- 10 quantification of closely related DNA or RNA sequences.

Methods which may also be used to quantify the expression of HRP include radiolabeling or biotinylating nucleotides, compaction of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplas, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be
- 15 accelerated by running the assay in a high-throughput format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

- Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g.,
- 25 Brenan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Scheina, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Balasubramanian et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

- In another embodiment of the invention, nucleic acid sequences encoding HRP may be used
- 30 to generate hybridization probes useful in mapping the naturally occurring genome sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent *in situ* hybridization (FISH) may be correlated with other physical, chromosome mapping techniques and genetic map data. (See, e.g., Heitz-Jirlich, et al. (1995) in Myers, supra, pp. 965-988.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the
- 5 location of the gene encoding HRP on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

- In situ* hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides
- 15 valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., axatax-telangectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among
- 20 normal, carrier, or affected individuals.

In another embodiment of the invention, HRP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes
- 25 between HRP and the agent being tested may be measured.

- Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Oryen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with HRP, or fragments thereof,
- 30 and washed. Bound HRP is then detected by methods well known in the art. Purified HRP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding HRP specifically compete with a test compound for binding HRP. In
- 35 this manner, antibodies can be used to detect the presence of any peptide which shares one or more

antigenic determinants with HRP.

In additional embodiments, the nucleotide sequences which encode HRP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the right genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative and not limitative of the remainder of the disclosure in any way whatsoever.

- 10 The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. 60/125,593; U.S. Ser. No. 60/133,040; and U.S. Ser. No. 60/143,188, are hereby expressly incorporated by reference.

EXAMPLES

1. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in lysis buffer or in a suitable mixture of denaturants, such as TRIzol (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl gradients or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with Dnase. For most libraries, poly(A⁺) RNA was isolated using oligo(dT)-coupled paramagnetic particles (Promege), OLIGOEX, Inc. particles (QIAGEN), Chatsworth, CA, or an OLIGOEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin, TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPRESCRIPT[®] plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo(dT) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHARCYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column

chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the phylphinder of a suitable plasmid, e.g.,

5 PBLUESCRIPT[®] plasmid (Stratagene), pSVKRT[®] plasmid (Life Technologies), pCDNA2.1 plasmid (Invitrogen), Cusbind[®] CA, or pINCY[®] plasmid (Incyte Pharmaceuticals, Palo Alto, CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueRF[®], or SOL[®] from Stratagene or DH5α, DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Miniprep DNA purification system (Promege), an ACTE Miniprep purification kit (Epic Biosystems, Gaithersburg, MD), and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

15 Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rea, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene, OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

cDNA sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYT[®] 800 (Pr4Fin-Ether) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microprocessor (BioBios Scientific) or the MICROLAB 200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIODYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Perkin-Elmer) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 77). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example V.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed

PCT/US00/07277

The concentration of DNA in each well was determined by dissolving 100 μ l of PICOgreen quantitation reagent (0.23% w/v) PICOgreen, Molecular Probes, Eugene, OR) into 100 μ l of 1 X TE and 0.5 μ l of unlabelled PCR product into each well of an opaque fluorescent plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a fluoroscan II (LabSystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analysed by electrophoresis on a 1% agarose mini-gel to determine which reactions were successful in extending the sequence.

The cells were lysed and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, 4, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYEANALYTIC energy transfer sequencing primers and the DYEANALYTIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIODYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

5 Hybridization probes (derived from SEQ ID NOs:1-58) are employed to screen cDNAs, genomic DNAs, or mRNAs. Alternatively, the labeling of oligonucleotide, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 μ mol of each oligo, 250 μ Ci of γ -³²P-labeled dATP (Amersham Pharmacia Biotech), and T₄ polynucleotide kinase (DDBP NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPRALABEX G-25 superfine size exclusion column (Amersham Pharmacia Biotech). An aliquot containing 10 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis. Human DNA digested with one or the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DDBP NEN).

A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Buehler, US2001/0011122A1)

23. do or did not may also be used to arrange and fix elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or

selected at random from a cDNA library relevant to the present invention, are arranged on appropriate substrates, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:339-345.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

VIII. Complementary Polynucleotides

Sequences complementary to the HRP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring HRP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of HRP. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the HRP-encoding transcription.

IX. Expression of HRIP

Expression and purification of HRP is achieved using bacterial or virus-based expression systems. For expression of HRP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *rnp-lac* (*lac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory elements. Recombinant viruses are transformed into suitable bacterial hosts, e.g., BL21 (DE3).

Antibiotic resistant bacteria express HRP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of HRP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographa californica nuclear polyhedrosis virus

to be known as baculovirus. The nonessential polyhedrin gene of baculovirus is (ACMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding HRP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculoviruses in some cases infect *Sitotrupa frugiperda* (SF9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engsthalder, E.K. et al. (1993a) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945).

affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kDa enzyme from *Saccharomyces glutathionyltransferase*, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Averbach Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from HRP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunofluorescence purification using commercially available monoclonal anti-FLAG antibodies (Kodak). His₆, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995), Sambrook et al., 10 and 16. Purified HRP obtained by these methods can be used directly in the following activity assay.

X. Demonstration of HRIP Activity

kinase activity of HRP is measured by the phosphorylation of appropriate substrates using gamma-labeled γ -ATP and quantitation of the incorporated radioactivity using a beta radiocounter. HRP is incubated with the protein substrate, γ -ATP, and an appropriate kinase buffer. The γ -ATP incorporated into the product is separated from free γ -ATP by electrophoresis and the incorporated γ is counted. The amount of γ recovered is proportional to the kinase activity of HRP in the assay. A determination of the specific amino acid residue phosphorylated by protein kinase activity is made by phosphoamino acid analysis of the hydrolyzed protein.

Alternatively, protein phosphatase activity of HRP is measured by the hydrolysis of *p*-nitrophenyl phosphate (PNPP). HRP is incubated together with PNPP in HEPES buffer pH 7.5, in the presence of 0.1% *m*-cresol/ethanol at 37°C for 60 min. The reaction is stopped by the addition of 6 ml of 10 N NaOH and the increase in light absorbance at 410 nm resulting from the hydrolysis of PNPP is measured using a spectrophotometer. The increase in light absorbance is proportional to the activity of HRP in the assay (Diamond, R.H. et al., 1994) (Mol. Cell Biol. 14:3753-3762).

25 **XI. Functional Assays**

HRP function is assessed by expressing the sequences encoding HRP in physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT plasmid (Life Technologies) and pCL3.1 plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5–10 µg of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1–2 µg of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP).

- Clonetics), CD44 or a CD44-GFP fusion protein. Flow cytometry (FACS), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD44-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ornsted, M.G. (1994) *Elsevier*, Oxford, New York NY.
- The influence of HRP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding HRP and either CD44 or CD44-GFP. CD44 and CD44-GFP are expressed on the surface of transfected cells and bind to connected regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD44 (DONALD, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding HRP and other genes of interest can be analyzed by northern analysis or microarray techniques.

XII. Production of HRP-Specific Antibodies

- HRP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M. G. (1990) *Methods Enzymol.* 182:483-493), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

- Alternatively, the HRP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophobic regions are well described in the art (See, e.g., Ausubel, 1995, *JMBA*, ch. 11).

- Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Pierce-Albany) using fine-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidebenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *JMBA*). Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-HRP activity by, for example, binding the peptide or HRP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-labeled goat anti-rabbit IgG.

XIII. Purification of Naturally Occurring HRP Using Specific Antibodies

- Naturally occurring or recombinant HRP is substantially purified by immunoaffinity chromatography using antibodies specific for HRP. An immunoaffinity column is constructed by covalently coupling anti-HRP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blebbed and washed according to the manufacturer's instructions.

- Media containing HRP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential abundance of HRP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/HRP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and HRP is collected.

XIV. Identification of Molecules Which Interact with HRP

- HRP, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539). Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled HRP, washed, and any wells with labeled HRP complex are assayed. Data obtained using different concentrations of HRP are used to calculate values for the number, affinity, and association of HRP with the candidate molecules.

- Alternatively, molecules interacting with HRP are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1988), *Nature* 340:245-246), or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

- Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described method for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1 (cont.)

Protein Seq ID	Accession ID	Clone ID	Library	Fragment(s)
10	U01481	1673761	BLADNOT05	167376166 (BLADNOT05) 202481566 (KBA1NOT05) 437861811 (BLADNOT22)
11	U120442	BLA1NOT05	40162986 (BLA1NOT05) 127044284 (BLA1NOT05) 167250666 (BLA1NOT05)	
12	U187713	LSK1NOT03	329368814 (LSP1NOT03) 296658111 (LSP1NOT03) 274661401 (LSP1NOT03)	
13	U187713	LSK1NOT03	296658114 (LSP1NOT03) 142714266 (K11NOT03) 187713281 (LSK1NOT03)	
14	U271615	THA1NOT04	26375966 (BOM1NOT04) 263759658 (BOM1NOT04) 07917481 (SYN1NOT04) 12721240281 (LST1NOT04) 12721240281 (LST1NOT04) 266802741 (THA1NOT04) 227295641 (THA1NOT04)	
15	U271615	THA1NOT04	26375966 (BOM1NOT04) 263759658 (BOM1NOT04) 07917481 (SYN1NOT04) 12721240281 (LST1NOT04) 12721240281 (LST1NOT04) 266802741 (THA1NOT04) 227295641 (THA1NOT04)	

Table 1

Seq ID	Seq ID No.	Accession No.	Library	Fragment
1	480457	151	LIBRARY	1547596 (TSPH024) 480457 (TSPH024) 18736866 (PROMOT02) 11489676 (ADENOM04) 37667441 (BRST024)
2	16	17	REFINOT01	256464X (BOS18E02) 205644X (BOS18E02) 666131X (HEMLP01)
3	1452425	17	REFINOT01	1207366X (MUSGNT01) 1475842X (REFINOT01) 15717293X (UTRSMOT05)
4	2439407	4	CONSACT01	1630939X (BRST064) 35400247X (BRST064) 15609097X (BRST064)
5	2451617	5	HNH1ZMT01	4742686X (H1A001) 9412941X (ADENOM03) 14219517X (HNH1ZMT01)
6	205186	20	TOMSMOT01	1759582X (T1UMOT03) 3015186X (TOMSMOT01) 584202766X (TOMSMOT01)
7	5504546	21	RHBDIN01	1759572X (TOMSMOT01) 12917847X (PROMOT03) 5504546X (RHBADIN01)
8	1511326	22	LUNSMOT1	1511326X (LUNSMOT1) 2922438X (LUNSMOT1) 1511326X (LUNSMOT1)
9	1519120	23	BLMDV004	1519120X (BLMDV004) 1519120X (BLMDV004) 1519120X (BLMDV004)

Table 3 (cont.)

Polynucleotide SEQ ID NO:	Useful Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
24	163-207	Nervous (0.133) Cardiovascular (0.133) Dermatologic (0.133)	Cancer (0.400) Fetal/Cell Line (0.200) Inflammation (0.133)	pINCY
25	19-63	Cardiovascular (0.182) Hematopoietic/Immune (0.182) Reproductive (0.182)	Cancer (0.515) Cell Proliferation (0.242) Inflammation (0.242)	pINCY
26	297-343	Cardiovascular (0.250) Hematopoietic/Immune (0.150) Musculoskeletal (0.150)	Cancer (0.300) Cell Proliferation (0.250) Inflammation (0.200)	pINCY
27	271-315	Endocrine (0.500) Musculoskeletal (0.500)	Cancer (0.500)	pINCY
28	161-207	Reproductive (0.241) Gastrointestinal (0.233) Cardiovascular (0.150)	Cancer (0.429) Inflammation (0.263) Cell Proliferation (0.211)	pINCY

Table 4

Polynucleotide SEQ ID NO:	Library	Library Comment
15	LIVBCT01	Library was constructed using RNA isolated from the liver tissue of a patient with primary biliary cirrhosis who had a liver transplant.
16	NEUTLPT01	Library was constructed using RNA isolated from peripheral blood granulocytes collected by density gradient centrifugation through Ficoll-Hypaque. The cells were isolated from buffy coat units obtained from unrelated male and female donors. Cells were cultured in 100 ng/ml E. coli LPS for 30 minutes.
17	BEPINON01	This normalized bronchial epithelium library was constructed from 5.12 million independent clones from a bronchial epithelium library. RNA was made from a bronchial epithelium primary cell line derived from a 54-year-old Caucasian male. The normalization and hybridization conditions were adapted from Soares et al., Proc. Natl. Acad. Sci. USA (1994) 91:9228, using a longer (24-hour) reannealing hybridization period.
18	COLSUCT01	Library was constructed using RNA isolated from diseased sigmoid colon tissue obtained from a 70-year-old Caucasian male during colectomy with permanent ileostomy. Pathology indicated chronic ulcerative colitis. Patient history included benign neoplasm of the colon. Family history included atherosclerotic coronary artery disease and myocardial infarctions.
19	HNTJAZT01	Library was constructed using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated for three days with 0.35 micromolar 5-aza-2'-deoxycytidine (AZ).
20	TONSNOT03	Library was constructed using RNA isolated from diseased left tonsil tissue removed from a 6-year-old Caucasian male during adenotonsillectomy. Pathology indicated reactive lymphoid hyperplasia, bilaterally. Family history included benign hypertension, myocardial infarction, and atherosclerotic coronary artery disease.
21	BRABDI01	Library was constructed using RNA isolated from diseased cerebellum tissue removed from the brain of a 57-year-old Caucasian male, who died from a cerebrovascular accident.
22	LUNGNOT14	Library was constructed using RNA isolated from lung tissue removed from the left lower lobe of a 47-year-old Caucasian male during a segmental lung resection. Pathology for the associated tumor tissue indicated a grade 4 adenocarcinoma, and the parenchyma showed calcified granuloma. Patient history included benign hypertension and chronic obstructive pulmonary disease. Family history included type II diabetes and acute myocardial infarction.

Polynucleotide 550 ID NO.	Library	Library Comment
23	BLADT004	Library was constructed using RNA isolated from bladder tumor tissue removed from a 60-year-old Caucasian male during a radical cystectomy, prostatectomy, and vasectomy. Pathology in-situ was identified in the dome and trigone. Family history included type 3 osteosarcoma, closed fracture, and type 1 diabetes.
24	BLADH005	Library was constructed using RNA isolated from bladder tissue removed from a 60-year-old Caucasian male during a radical cystectomy, prostatectomy, and vasectomy. Pathology in-situ was identified in the dome and trigone. Family history included type 3 osteosarcoma, closed fracture, and type 1 diabetes.
25	BLAH009	Library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus, who died at 23 weeks gestation.
26	LEUKH003	Library was constructed using RNA isolated from white blood cells of 27-year-old female with blood type A-. The donor tested negative for cytomegalovirus (CMV).
27	BONM001	Library was constructed using RNA isolated from clonal peritoneum removed from a 20-year-old Caucasian male during a laparoscopy with inspection above the nore. Pathology for the associated tumor tissue indicated partially necrotic and cystic osteoblastic grade 3 osteosarcoma (post-chemotherapy). Family history included osteogenesis imperfecta, closed fracture, and type 1 diabetes.
28	THYRH009	Library was constructed using RNA isolated from diseased thyroid tissue removed from an 18-year-old Caucasian female during an unilateral thyroid lobectomy and regional lymph node excision. Pathology indicated adenomatous nodule. This was associated with a follicular adenoma of the thyroid. Family history included thyroid cancer.

Table 4 (cont.)

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Foster City, CA. Foster-Eimer Applied Biosystems.	
ABI PARACEL PDF	A Fast Plus Finder used in comparing and annotating amino acid or nucleic acid sequences.	Foster City, CA; Packard Inc., Pasadena, CA.	Mismatch <50%
ABI AnnuAsubmitter	A program that assembles nucleic acid sequences.	Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST finds five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990). J. Mol. Biol. 215: 403-410. Altschul, S.F. et al. (1997). Nucleic Acids Res. 25: 3389-3402.	EST: Probability value 1.0E-8 or values 1.0E-10 or less EST: E value 1.0E-6 ATestbed EST: E value 1.0E-6 95% or greater and Match length > 200 bases or greater: blastx E value 1.0E-8 or less blastn E value 1.0E-10 or greater blastx E value 1.0E-8 or less blastn E value 1.0E-10 or greater
PASTA	A Pairwise Alignment Searcher that matches a sequence similarity between a query sequence and a group of sequences of the same type. PASTA compares at least five functions: blastp, blastn, blastx, tblastn, and tblastx.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1996) Methods Enzymol. 183: 63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	Score 1000 or greater: Ratio of Score/Length = 0.75 or greater and 10 applicable. Probability value 1.0E-3 or less Score 10 50 bits for PFAM hits, depending on individual protein families
BLIMP5	A Blocks Improved Searcher that matches a sequence homology, and structural fingerprint regions.	Henikoff, S. and J.C. Henikoff, Nucleic Acids Res. 19:655-72, 1991. J.C. Henikoff and Henikoff (1996) Methods Enzymol. 266: 105-132; and Altschul, S.F. et al. (1997). J. Chem. Inf. Comput. Sci. 37: 417-424.	Score 10 50 bits for PFAM hits, depending on individual protein families
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM) based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994). J. Mol. Biol. 233:1301-1331; Sommeringer, E.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score 10 50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
PrositeScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribkov, M. et al. (1988) CABIOS 4:61-66; Gribkov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Normalized quality score; GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and Crossmatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
CuSed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:193-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=3.5 or greater
Mutis	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <i>supra</i> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

WFO 0605332

PCT/US98/07177

WFO 0805

PCT/US98/07177

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:
 - a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-14,
 - b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-14,
 - c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-14, and
 - d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-14.
2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-14.
3. An isolated polynucleotide encoding a polypeptide of claim 1.
4. An isolated polynucleotide of claim 3 selected from the group consisting of SEQ ID NO:15-28.
5. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
6. A cell transformed with a recombinant polynucleotide of claim 5.
7. A transgenic organism comprising a recombinant polynucleotide of claim 5.
8. A method for producing a polypeptide of claim 1, the method comprising:
 - a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and
 - b) recovering the polypeptide so expressed.
9. An isolated antibody which specifically binds to a polypeptide of claim 1.

10. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:
- a polynucleotide sequence selected from the group consisting of SEQ ID NO:15-28;
 - a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:15-28;
 - a polynucleotide sequence complementary to a),
 - a polynucleotide sequence complementary to b), and
 - an RNA equivalent of a)-d).
11. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 10.
12. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 10, the method comprising:
- hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and
 - detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.
13. A method of claim 12, wherein the probe comprises at least 30 contiguous nucleotides.
14. A method of claim 12, wherein the probe comprises at least 60 contiguous nucleotides.
15. A pharmaceutical composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.
16. A method for treating a disease or condition associated with decreased expression of functional HRIP, comprising administering to a patient in need of such treatment the pharmaceutical composition of claim 15.
17. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:
- exposing a sample comprising a polypeptide of claim 1 to a compound, and

- detecting agonist activity in the sample.

18. A pharmaceutical composition comprising an agonist compound identified by a method of claim 17 and a pharmaceutically acceptable excipient.

19. A method for treating a disease or condition associated with decreased expression of functional HRIP, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 18.

20. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- exposing a sample comprising a polypeptide of claim 1 to a compound, and
- detecting antagonist activity in the sample.

21. A pharmaceutical composition comprising an antagonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

22. A method for treating a disease or condition associated with overexpression of functional HRIP, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 21.

23. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 4, the method comprising:

- exposing a sample comprising the target polynucleotide to a compound, and
- detecting altered expression of the target polynucleotide.

PF-0683 PCT

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.

BANDMAN, Olga

TAN, Y. Tom

HILMAN, Jennifer L.

BANCRO, Mariah R.

AZIMZAI, Yalda

LU, Dongyuan Aina H.

MO-TOHOK, Janice

<120> REGULATORS OF INTRACELLULAR PHOSPHORYLATION

<130> PF-0683 PCT

<140> To Be Assigned

<151> Herewith

<150> 60/125,593, 60/135,049, 60/143,188
<152> 1999-03-18, 1999-05-20, 1999-07-09

<160> 28

<170> FEEL Program

<210> 1

<211> 482

<212> PRT

<213> Homo sapiens

<220>

<221> misc:feature

<223> Inocyte ID No: 480657CD1

<400> 1

Met Pro Ser Pro Leu Asp Asp Arg Val Val Ala Leu Ser

1 10 15

Arg Pro Val Arg Pro Asp Leu Asn 25 30

Tyr Leu Gly Ser Ala Asn Ser His Pro Pro Val

35 40 45

Ile Ala Thr Thr Val Ser Leu Lys Ala Ala Asn Leu Thr Tyr

Met Pro Ser Ser Ser Cys Ser Leu Asn Cys Gly Cys

65 70 75

Ser Ser Ala Ser Cys Cys Thr Val Ala Thr Tyr Asp Lys Asp Asn

80 85 90

Gln Ala Gln Thr Gln Gly Thr Thr Thr Thr Thr Thr Thr Thr

95 100 105

Ile Gly Thr Ser Thr Thr Pro Ala Asn Gln Met Val Asn Asn

110 115 120

Asn Gln Asn Thr Gly Ser Leu Ser Pro Ser Cys Gly Val Gly

125 130 135

Pro Val Ser Gly Thr Pro Lys Gln Leu Ala Ser Ile Lys Ile

140 145 150

Tyr Pro Asn Asp Leu Ala Lys Lys Met Thr Lys Cys Ser Lys Ser

155 160 165

His Leu Pro Ser Thr Gly Pro Val Ile Ile Asp Cys Arg Pro Phe

170 175 180

PF-0683 PCT

Met Gly Tyr Asn Lys Ser His Ile Gln Gly Ala Val His Ile Asn

185 190 195

Cys Ala Asp Lys Ile Ser Arg Arg Leu Gln Gln Gly Lys Ile

200 205 210

Thr Val Leu Asp Leu Ile Ser Cys Arg Gly Cys Lys Asp Ser Phe

215 220 225

Lys Arg Ile Phe Ser Lys Gln Ile Ile Val Tyr Asp Gln Asn Thr

230 235 240

Asn Gln Pro Ser Thr Val Met Pro Leu His Ile Gln

245 250 255

Leu Gly Ser Leu Lys Arg Gly Lys Gly Pro Leu Val Leu Lys

260 265 270

Gly Gly Leu Ser Ser Phe Lys Gln Asn His Gln Asn Leu Cys Asp

275 280 285

Asn Ser Leu Gln Leu Gln Gly Arg Gln Val Gly Gly Cys Ala

290 295 300

Ser Ala Ala Ser Ser Ser Leu Pro Gln Pro Ile Pro Thr Phe

305 310 315

Asp Ile Gln Asn Gln Thr Pro Lys Pro Lys Pro Gln Leu

320 325 330

Leu Gly Asn Gln Gln Asp Ala Gln Asp Leu Asp Thr Met Gln Arg

335 340 345

Leu Asn Ile Gly Tyr Val Ile Asn Val Thr Thr His Leu Pro Asn

350 355 360

Tyr His Tyr Gln Lys Gly Leu Phe Asn Tyr Lys Arg Leu Pro Ala

365 370 375

Thr Asp Ser Asn Lys Gln Asn Leu Arg Gln Tyr Phe Gln Gln Ala

380 385 390

Phe Gln Phe Ile Gln Gln Ala His Gln Cys Gly Lys Gly Leu Leu

395 400 405

Ile His Cys Gln Ala Gly Val Ser Arg Ser Ala Thr Ile Val Ile

410 415 420

Ala Tyr Leu Met Cys His Thr Arg Met Thr Met Thr Asp Ala

425 430 435

Lys Phe Val Lys Gly Lys Arg Pro Ile Ile Ser Pro Asn Leu Asn

440 445 450

Phe Met Gly Gln Leu Leu Gln Phe Gln Gln Asp Leu Asn Asn

455 460 465

Val Thr Pro Arg Ile Leu Thr Pro Lys Leu Met Gly Val Gln Thr

470 475 480

Val Val

<210> 2

<211> 190

<212> PRT

<213> Homo sapiens

<220>

<221> misc:feature

<223> Inocyte ID No: 50565CD1

<400> 2

Met Ser Arg Arg Arg Phe Asp Cys Arg Ser Ile Ser Gly Leu Leu

1 5 10 15

Thr Thr Thr Pro Gln Ile Pro Ile Lys Met Gln Asn Phe Asn Asn

16 20 25

Phe Tyr Ile Leu Thr Ser Lys Gln Leu Gly Arg Gly Lys Phe Ala

26 30 35 40 45

Val Val Arg Gln Gln Cys Ile Ser Lys Ser Thr Gly Gln Gln Tyr Ala

50 55

PF-0683 NCT

Gly Ser Gln Ser Gly Thr Gly Ser Gly Gln Pro Gly Gly Thr Leu
140 145 150 155 160 165 170 175 180 185 190 195 200 205 210
Gly Tyr Leu Ala 140 145 150 155 160 165 170 175 180 185 190 195 200 205 210
Ser Thr Ala Ser Asp Val Tyr Ser Phe Gly Ile Leu Met Trp Ala
170 175 180 185 190 195 200 205 210 215 220 225 230 235 240
Val Leu Ala Gly Asp Gln Val Gln Leu 170 175 180 185 190 195 200 205 210 215 220 225 230 235 240
Val Tyr Gln Ala Val Cys Asn Arg Gln Asn Arg Pro Ser Leu Ala
200 205 210 215 220 225 230 235 240 245 250 255 260 265 270
Glu Leu Pro Gln Ala Gly Pro Gln Thr Pro Gly Leu Gln Gly Leu
210 215 220 225 230 235 240 245 250 255 260 265 270 275 280
Lys Gln Leu Met Gln Leu Cys Trp Ser Ser Gln Pro Lys Asp
230 235 240 245 250 255 260 265 270 275 280 285 290 295 300
Pro Ser Phe Gln Gln Cys Leu Pro Lys Thr Asp Gln Val Phe Gln
240 245 250 255 260 265 270 275 280 285 290 295 300 305 310
Met Val Gln Asn Asn Met Asn Ala Val Ser Thr Val Lys Asp
260 265 270 275 280 285 290 295 300 305 310 315 320 325 330
Phe Leu Ser Gln Leu Arg Ser Asn Arg Asp Phe Ser Ile Pro
270 275 280 285 290 295 300 305 310 315 320 325 330 335 340
Glu Ser Gln Gly Gly Thr Gln Met Asp Gly Phe Arg Arg 300
280 285 290 295 300 305 310 315 320 325 330 335 340 345 350
Ile Gln Asn Gln His Ser Arg Asn Asp Val Met Ser Val Pro Lys
300 305 310 315 320 325 330 335 340 345 350 355 360 365 370
Leu Asn Lys Leu 300 305 310 315 320 325 330 335 340 345 350 355 360 365 370
Lys Cys Pro Ser Leu Thr Lys Arg Ser Arg Ala Gln Gln Gly Trp
310 315 320 325 330 335 340 345 350 355 360 365 370 375 380
Val Pro Gln Ala 310 315 320 325 330 335 340 345 350 355 360 365 370 375 380
Gln Pro Pro Gln Thr Pro Gln Thr Ser Thr Phe Arg Asn Gln Met
340 345 350 355 360 365 370 375 380 385 390 395 400 405 410
Pro Ser Pro Thr 380 385 390 395 400 405 410 415 420 425 430 435 440
Asn Gln Gly Ala Gln Arg Gln Gly Met Asn Trp Ser Cys Arg Thr
390 395 400 405 410 415 420 425 430 435 440 445 450 455 460
Pro Gln Pro Asn Pro Val Thr Gly Arg 410 415 420 425 430 435 440 445 450 455 460 465 470
Asn Cys Ser Gly Val Gln Val Gly Asp Asn Thr Leu Thr Met
425 430 435 440 445 450 455 460 465 470 475 480 485 490 495 500
Gln Gln Thr Thr Ala Leu Pro Thr Trp Gly Leu Ala Pro Ser Gly
435 440 445 450 455 460 465 470 475 480 485 490 495 500 505 510
Lys Gly Arg Gly Leu Gln His Pro Pro Val Gly Ser Gln Gly
455 460 465 470 475 480 485 490 495 500 505 510 515 520 525 530
Gly Pro Lys Asp Pro Gln Ala Trp Ser Arg Pro Gln Gly Tyr Trp
485 490 495 500 505 510 515 520 525 530 535 540 545 550 555 560
Asn His Ser Gly Lys 485 490 495 500 505 510 515 520 525 530 535 540 545 550 555 560

<210> 5
<211> 384
<212> PRT
<213> Homo sapiens
<220>
<221> misc-feature
<223> InKey ID No: 34156170D1
<400> 5
Met Asp Pro Ala Gly Pro Arg Gly Val Leu Pro Arg Pro Cys

5/24

PF-0683 PCT

Arg Val Leu Val Leu Leu Asn Pro Arg Gln Gly Lys Gly Lys 15
1 5 10 15
Leu Gln Leu Phe Arg Ser His Val Gln Pro Leu Leu Ala Gln Ala 20 25 30 35 40 45
Glu Ile Ser Phe Thr Met Leu Thr Gln Arg Arg Asn His Ala 35 40 45 50 55 60
Arg Gln Leu Val Arg Ser Gln Val Met Gly Arg Trp Asp Ala Leu 65 70 75 80 85 90
Val Val Met Ser Gly Asp Gly Leu Met His Gln Val Val Asn Gly 65 70 75 80 85 90
Leu Met Gln Arg Pro Asp Trp Gln Thr Ala Ile Gln Lys Pro Leu 95 100 105 110 115 120
Cys Ser Leu Pro Ala Gly Ser Gly Asn Ala Leu Ala Ala Ser Leu 95 100 105 110 115 120
Asn His Tyr Ala Gly Tyr Gln Gln Val Thr Asn Gln Asp Leu 115 120 125 130 135 140
Thr Asn Cys Thr Leu Leu Cys Arg Arg Asp Leu Ser Pro Met 120 125 130 135 140 145
Asn Leu Leu Ser 135 140 145 150 155 160 165 170 175 180 185 190 195 200 205 210
Val Leu Ser Leu Ala Trp Gly Phe Ile Ala Asp Val Asp Leu Gln 140 145 150 155 160 165 170 175 180 185 190 195 200 205 210
Ser Gln Lys Tyr 185 190 195 200 205 210 215 220 225 230 235 240 245 250 255 260
Thr Phe Leu Arg Leu Ala Leu Arg Thr Tyr Arg Gly Arg Leu 210 215 220 225 230 235 240 245 250 255 260 265 270
Ala Tyr Leu Pro 200 205 210 215 220 225 230 235 240 245 250 255 260 265 270
Pro Val Val Val Gln Gln Gly Pro Val Asp Ala His Leu Val Pro 240 245 250 255 260 265 270 275 280 285 290 295 300 305 310
Leu Gln Gln Pro 230 235 240 245 250 255 260 265 270 275 280 285 290 295 300 305 310
Lys Asp Phe Val Ala Pro Met Gly Arg Cys Ala Ala Gly Val Met 260 265 270 275 280 285 290 295 300 305 310 315 320 325 330 335 340
His Leu Phe Tyr Val Arg Ala Gly Val Ser Arg Ala Met Leu 300 305 310 315 320 325 330 335 340 345 350 355 360 365 370 375 380
Arg Leu Phe Leu Ala Met Gln Lys Gly Arg His Met Gln Tyr Gln 290 295 300 305 310 315 320 325 330 335 340 345 350 355 360 365 370 375 380
Cys Pro Tyr Leu Val Tyr Val Pro Val 320 325 330 335 340 345 350 355 360 365 370 375 380 385 390 395 400 405 410
Pro Lys Asp Gly Lys Gly Val Phe Ala Val Asp Gly Gln Leu Gln 320 325 330 335 340 345 350 355 360 365 370 375 380 385 390 395 400 405 410
Val Ser Gln Ala 330 335 340 345 350 355 360 365 370 375 380 385 390 395 400 405 410
Met Val Ser Gly Cys Val Gln Pro Pro Ser Trp Lys Pro Gln 350 355 360 365 370 375 380 385 390 395 400 405 410 415 420 425 430 435 440
Gln Met Pro Pro 380 385 390 395 400 405 410 415 420 425 430 435 440 445 450 455 460 465 470 475 480 485 490 495 500 505 510 515 520 525 530 535 540 545 550 555 560 565 570 575 580 585 590 595 600 605 610 615 620 625 630 635 640 645 650 655 660 665 670 675 680 685 690 695 700 705 710 715 720 725 730 735 740 745 750 755 760 765 770 775 780 785 790 795 800 805 810 815 820 825 830 835 840 845 850 855 860 865 870 875 880 885 890 895 900 905 910 915 920 925 930 935 940 945 950 955 960 965 970 975 980 985 990 995 1000 1005 1010 1015 1020 1025 1030 1035 1040 1045 1050 1055 1060 1065 1070 1075 1080 1085 1090 1095 1100 1105 1110 1115 1120 1125 1130 1135 1140 1145 1150 1155 1160 1165 1170 1175 1180 1185 1190 1195 1200 1205 1210 1215 1220 1225 1230 1235 1240 1245 1250 1255 1260 1265 1270 1275 1280 1285 1290 1295 1300 1305 1310 1315 1320 1325 1330 1335 1340 1345 1350 1355 1360 1365 1370 1375 1380 1385 1390 1395 1400 1405 1410 1415 1420 1425 1430 1435 1440 1445 1450 1455 1460 1465 1470 1475 1480 1485 1490 1495 1500 1505 1510 1515 1520 1525 1530 1535 1540 1545 1550 1555 1560 1565 1570 1575 1580 1585 1590 1595 1600 1605 1610 1615 1620 1625 1630 1635 1640 1645 1650 1655 1660 1665 1670 1675 1680 1685 1690 1695 1700 1705 1710 1715 1720 1725 1730 1735 1740 1745 1750 1755 1760 1765 1770 1775 1780 1785 1790 1795 1800 1805 1810 1815 1820 1825 1830 1835 1840 1845 1850 1855 1860 1865 1870 1875 1880 1885 1890 1895 1900 1905 1910 1915 1920 1925 1930 1935 1940 1945 1950 1955 1960 1965 1970 1975 1980 1985 1990 1995 2000 2005 2010 2015 2020 2025 2030 2035 2040 2045 2050 2055 2060 2065 2070 2075 2080 2085 2090 2095 2100 2105 2110 2115 2120 2125 2130 2135 2140 2145 2150 2155 2160 2165 2170 2175 2180 2185 2190 2195 2200 2205 2210 2215 2220 2225 2230 2235 2240 2245 2250 2255 2260 2265 2270 2275 2280 2285 2290 2295 2300 2305 2310 2315 2320 2325 2330 2335 2340 2345 2350 2355 2360 2365 2370 2375 2380 2385 2390 2395 2400 2405 2410 2415 2420 2425 2430 2435 2440 2445 2450 2455 2460 2465 2470 2475 2480 2485 2490 2495 2500 2505 2510 2515 2520 2525 2530 2535 2540 2545 2550 2555 2560 2565 2570 2575 2580 2585 2590 2595 2600 2605 2610 2615 2620 2625 2630 2635 2640 2645 2650 2655 2660 2665 2670 2675 2680 2685 2690 2695 2700 2705 2710 2715 2720 2725 2730 2735 2740 2745 2750 2755 2760 2765 2770 2775 2780 2785 2790 2795 2800 2805 2810 2815 2820 2825 2830 2835 2840 2845 2850 2855 2860 2865 2870 2875 2880 2885 2890 2895 2900 2905 2910 2915 2920 2925 2930 2935 2940 2945 2950 2955 2960 2965 2970 2975 2980 2985 2990 2995 3000 3005 3010 3015 3020 3025 3030 3035 3040 3045 3050 3055 3060 3065 3070 3075 3080 3085 3090 3095 3100 3105 3110 3115 3120 3125 3130 3135 3140 3145 3150 3155 3160 3165 3170 3175 3180 3185 3190 3195 3200 3205 3210 3215 3220 3225 3230 3235 3240 3245 3250 3255 3260 3265 3270 3275 3280 3285 3290 3295 3300 3305 3310 3315 3320 3325 3330 3335 3340 3345 3350 3355 3360 3365 3370 3375 3380 3385 3390 3395 3400 3405 3410 3415 3420 3425 3430 3435 3440 3445 3450 3455 3460 3465 3470 3475 3480 3485 3490 3495 3500 3505 3510 3515 3520 3525 3530 3535 3540 3545 3550 3555 3560 3565 3570 3575 3580 3585 3590 3595 3600 3605 3610 3615 3620 3625 3630 3635 3640 3645 3650 3655 3660 3665 3670 3675 3680 3685 3690 3695 3700 3705 3710 3715 3720 3725 3730 3735 3740 3745 3750 3755 3760 3765 3770 3775 3780 3785 3790 3795 3800 3805 3810 3815 3820 3825 3830 3835 3840 3845 3850 3855 3860 3865 3870 3875 3880 3885 3890 3895 3900 3905 3910 3915 3920 3925 3930 3935 3940 3945 3950 3955 3960 3965 3970 3975 3980 3985 3990 3995 4000 4005 4010 4015 4020 4025 4030 4035 4040 4045 4050 4055 4060 4065 4070 4075 4080 4085 4090 4095 4100 4105 4110 4115 4120 4125 4130 4135 4140 4145 4150 4155 4160 4165 4170 4175 4180 4185 4190 4195 4200 4205 4210 4215 4220 4225 4230 4235 4240 4245 4250 4255 4260 4265 4270 4275 4280 4285 4290 4295 4300 4305 4310 4315 4320 4325 4330 4335 4340 4345 4350 4355 4360 4365 4370 4375 4380 4385 4390 4395 4400 4405 4410 4415 4420 4425 4430 4435 4440 4445 4450 4455 4460 4465 4470 4475 4480 4485 4490 4495 4500 4505 4510 4515 4520 4525 4530 4535 4540 4545 4550 4555 4560 4565 4570 4575 4580 4585 4590 4595 4600 4605 4610 4615 4620 4625 4630 4635 4640 4645 4650 4655 4660 4665 4670 4675 4680 4685 4690 4695 4700 4705 4710 4715 4720 4725 4730 4735 4740 4745 4750 4755 4760 4765 4770 4775 4780 4785 4790 4795 4800 4805 4810 4815 4820 4825 4830 4835 4840 4845 4850 4855 4860 4865 4870 4875 4880 4885 4890 4895 4900 4905 4910 4915 4920 4925 4930 4935 4940 4945 4950 4955 4960 4965 4970 4975 4980 4985 4990 4995 5000 5005 5010 5015 5020 5025 5030 5035 5040 5045 5050 5055 5060 5065 5070 5075 5080 5085 5090 5095 5100 5105 5110 5115 5120 5125 5130 5135 5140 5145 5150 5155 5160 5165 5170 5175 5180 5185 5190 5195 5200 5205 5210 5215 5220 5225 5230 5235 5240 5245 5250 5255 5260 5265 5270 5275 5280 5285 5290 5295 5300 5305 5310 5315 5320 5325 5330 5335 5340 5345 5350 5355 5360 5365 5370 5375 5380 5385 5390 5395 5400 5405 5410 5415 5420 5425 5430 5435 5440 5445 5450 5455 5460 5465 5470 5475 5480 5485 5490 5495 5500 5505 5510 5515 5520 5525 5530 5535 5540 5545 5550 5555 5560 5565 5570 5575 5580 5585 5590 5595 5600 5605 5610 5615 5620 5625 5630 5635 5640 5645 5650 5655 5660 5665 5670 5675 5680 5685 5690 5695 5700 5705 5710 5715 5720 5725 5730 5735 5740 5745 5750 5755 5760 5765 5770 5775 5780 5785 5790 5795 5800 5805 5810 5815 5820 5825 5830 5835 5840 5845 5850 5855 5860 5865 5870 5875 5880 5885 5890 5895 5900 5905 5910 5915 5920 5925 5930 5935 5940 5945 5950 5955 5960 5965 5970 5975 5980 5985 5990 5995 6000 6005 6010 6015 6020 6025 6030 6035 6040 6045 6050 6055 6060 6065 6070 6075 6080 6085 6090 6095 6100 6105 6110 6115 6120 6125 6130 6135 6140 6145 6150 6155 6160 6165 6170 6175 6180 6185 6190 6195 6200 6205 6210 6215 6220 6225 6230 6235 6240 6245 6250 6255 6260 6265 6270 6275 6280 6285 6290 6295 6300 6305 6310 6315 6320 6325 6330 6335 6340 6345 6350 6355 6360 6365 6370 6375 6380 6385 6390 6395 6400 6405 6410 6415 6420 6425 6430 6435 6440 6445 6450 6455 6460 6465 6470 6475 6480 6485 6490 6495 6500 6505 6510 6515 6520 6525 6530 6535 6540 6545 6550 6555 6560 6565 6570 6575 6580 6585 6590 6595 6600 6605 6610 6615 6620 6625 6630 6635 6640 6645 6650 6655 6660 6665 6670 6675 6680 6685 6690 6695 6700 6705 6710 6715 6720 6725 6730 6735 6740 6745 6750 6755 6760 6765 6770 6775 6780 6785 6790 6795 6800 6805 6810 6815 6820 6825 6830 6835 6840 6845 6850 6855 6860 6865 6870 6875 6880 6885 6890 6895 6900 6905 6910 6915 6920 6925 6930 6935 6940 6945 6950 6955 6960 6965 6970 6975 6980 6985 6990 6995 7000 7005 7010 7015 7020 7025 7030 7035 7040 7045 7050 7055 7060 7065 7070 7075 7080 7085 7090 7095 7100 7105 7110 7115 7120 7125 7130 7135 7140 7145 7150 7155 7160 7165 7170 7175 7180 7185 7190 7195 7200 7205 7210 7215 7220 7225 7230 7235 7240 7245 7250 7255 7260 7265 7270 7275 7280 7285 7290 7295 7300 7305 7310 7315 7320 7325 7330 7335 7340 7345 7350 7355 7360 7365 7370 7375 7380 7385 7390 7395 7400 7405 7410 7415 7420 7425 7430 7435 7440 7445 7450 7455 7460 7465 7470 7475 7480 7485 7490 7495 7500 7505 7510 7515 7520 7525 7530 7535 7540 7545 7550 7555 7560 7565 7570 7575 7580 7585 7590 7595 7600 7605 7610 7615 7620 7625 7630 7635 7640 7645 7650 7655 7660 7665 7670 7675 7680 7685 7690 7695 7700 7705 7710 7715 7720 7725 7730 7735 7740 7745 7750 7755 7760 7765 7770 7775 7780 7785 7790 7795 7800 7805 7810 7815 7820 7825 7830 7835 7840 7845 7850 7855 7860 7865 7870 7875 7880 7885 7890 7895 7900 7905 7910 7915 7920 7925 7930 7935 7940 7945 7950 7955 7960 7965 7970 7975 7980 7985 7990 7995 8000 8005 8010 8015 8020 8025 8030 8035 8040 8045 8050 8055 8060 8065 8070 8075 8080 8085 8090 8095 8100 8105 8110 8115 8120 8125 8130 8135 8140 8145 8150 8155 8160 8165 8170 8175 8180 8185 8190 8195 8200 8205 8210 8215 8220 8225 8230 8235 8240 8245 8250 8255 8260 8265 8270 8275 8280 8285 8290 8295 8300 8305 8310 8315 8320 8325 8330 8335 8340 8345 8350 8355 8360 8365 8370 8375 8380 8385 8390 8395 8400 8405 8410 8415 8420 8425 8430 8435 8440 8445 8450 8455 8460 8465 8470 8475 8480 8485 8490 8495 8500 8505 8510 8515 8520 8525 8530 8535 8540 8545 8550 8555 8560 8565 8570 8575 8580 8585 8590 8595 8600 8605 8610 8615 8620 8625 8630 8635 8640 8645 8650 8655 8660 8665 8670 8675 8680 8685 8690 8695 8700 8705 8710 8715 8720 8725 8730 8735 8740 8745 8750 8755 8760 8765 8770 8775 8780 8785 8790 8795 8800 8805 8810 8815 8820 8825 8830 8835 8840 8845 8850 8855 8860 8865 8870 8875 8880 8885 8890 8895 8900 8905 8910 8915 8920 8925 8930 8935 8940 8945 8950 8955 8960 8965 8970 897

WO 00/55332

PF-0653 PCT

<400> 6
 Met Arg Tyr Tyr Gln Pro Pro Asp Asp Ile Leu Val Leu
 1 5 15
 Cys Leu Ser Tyr Ala Val Leu Met 25 30
 Gln Arg Asp Lys Arg Asp Thr Ser An Phe Asp Lys Glu Thr
 35 40 45
 Arg Gln Pro Val Gln Leu Thr Pro Thr Asp Lys Leu Phe Ile Met
 50 55
 Asn Leu Asp Gln An Gln Phe Ala Gly Phe Ser Tyr Thr An Pro
 60 65 70
 Glu Phe Val Ile 80

<210> 7

<211> 71

<212> PRT

<213> Homo sapiens

<220>

<221> misc:feature

<222> InChI ID No: 550454CDI

<400> 7

Met Leu Phe Gly Leu Val Arg Gln Gly Leu Lys Cys Asp Gly Cys
 1 5 10 15
 Gly Leu An Tyr His Lys Arg Cys Ala Phe Ser Ile Pro An An
 20 25 30
 Cys Ser Gly Ala Arg Lys Arg Arg Leu Ser Ile Pro An An
 35 40 45
 Ser Gly His Ser Val Arg Leu Gly Thr Ser Glu Ser Leu Pro Cys
 50 55
 Thr Ala Glu Leu Ser Arg Ser Thr Thr Thr Thr Thr Thr Thr
 60 65 70
 Arg Pro Ser Ser Ser Ser Ser Ala Ser Ser Tyr Tyr
 75
 Gly Arg Pro Ile 80
 Arg Arg Lys Met 85
 Arg Arg Lys Met 90
 Val Pro His Thr Phe Leu Ile His Ser Tyr Thr Arg Pro Thr Val
 95 100 105
 Cys Gln His Thr 110
 Lys Leu Lys 115
 Leu Phe Arg Gln 120
 Cys Gln Cys Lys Asp Cys Phe An Cys His Lys Arg Cys Ala
 125 130 135
 Thr Arg Val Pro 140
 An Cys Leu Gly 145
 Asp Val Pro Met Glu Glu Ala Thr Phe Ser Glu Ala Ser Lys
 150 155
 Ser Ala Leu Met 160
 Glu Ser Glu An 165
 Ser His Ser Glu An Ala His Ala Ser Glu Glu Gly
 170 175
 Glu Gly Gly Lys 180
 Gln Ser Ser Leu 185
 Tyr Ile Pro Leu Met 190
 Arg Val Val Gln Ser Val Arg His Thr Arg Lys Ser Thr
 195 200
 Thr Leu Arg Glu Thr Val His Thr Ser An Lys Asp Thr
 205 210
 Leu Arg Lys His Tyr Trp Arg Leu Asp Cys Lys Cys Ile Thr
 215 220
 225

7/28

WO 00/55332

PF-0653 PCT

Leu Phe Gln An Asn Thr Thr Asn Arg Tyr Tyr Lys Glu Ile Pro
 275 280
 Leu Ser Glu Ile 285
 Val Pro Gly Thr An Pro His Cys Phe Glu Ile Val Thr Ala
 290 295
 Asn Ala Thr Tyr 300
 Val Val Gly Glu Met 305
 Gly Pro Ser Gln Gly Ala Glu Ala Arg Gly Trp Glu Thr
 310 315
 Ala Ile Arg Gln 320
 Leu Met Pro Val 325
 Ser Ala Pro Gly His Ala Pro His Arg Gln Ala Ser Leu Asp Ala 330
 335
 Ser Val Ser An 340
 Gln Ile Gln Glu 345
 Val Tyr Gln Ile Phe Pro Asp Glu Val Leu Gly Ser Gly Gln Phe
 350 355
 Gly Val Tyr 360
 Gly Lys His Arg 365
 Ala Val Lys Val Ile Asp Lys Leu Arg Phe Pro Thr Lys Gln Glu
 370 375
 Ser Gln Leu Arg 380
 Val Phe Val Met Glu Lys Leu His Gly Asp Met Leu Glu Met
 385 390
 Pro Gly Ile Val An Leu Glu Cys Met Phe Glu Thr Pro Glu Lys
 395 400
 Ile Leu Ser Glu Lys Arg Leu Pro Glu Arg Leu Thr Lys
 405 410
 Phe Ile Thr Gln Ile Leu Val Ala Arg His Leu His Phe
 415 420
 Lys An Ile Val His Cys Asp Leu Lys Pro Glu An Val Leu
 425 430
 Ala Ser Ala Asp Pro Phe Pro Gln Val Lys Lys Asp Phe Gly
 435 440
 Phe Ala Arg Ile Ile Gly Glu Lys Ser 445
 Gly Thr Ala Tyr Leu Ala Pro Glu Val Leu An Gln Gly
 450 455
 Tyr An Arg Ser 460
 Asp Met Trp Ser 465
 Val Ser Leu Ser Gly Thr Phe Pro An Asp Glu Asp Ile
 470 475
 An Asp Gln Ile An Ala Phe 480
 Tyr Pro Ala Ser 485
 Trp Ser His Ser Ala Gly Ala Ile Asp Leu Ile An An Leu
 490 495
 Leu Gln Val Lys Arg Lys Arg Tyr Val Asp Lys Ser 500
 Ser His Pro Trp Leu Gln Tyr Gln Thr Leu Asp Leu Arg
 505 510
 Glu Leu Gly 515
 Met Gly Glu Arg 520
 Ile Thr His Glu 525
 Asp Asp Ala Arg Trp Glu Phe Ala Glu His Pro Leu Pro
 530 535
 Ser Gly Leu 540
 Thr Asp Arg Asp 545
 Pro Gln Asp His 550
 Asp Met Gln Ala Glu Arg Ile Ser Val
 555 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 Leu

8/24

WO 0065333

PCT/US00/0777

PF-0683 PCT

<210> 8
<211> 249
<212> PRT
<213> Homo sapiens
<220>
<221> misc_Feature
<222> Inocyte ID No: 151133C01
<400> 8
Met Ala Ser Ser Asp Asp Asp Gly Thr Arg Gly Ala Ser Gln
1 Ala Gly Gln Asp Val Gln Ala Pro Gly Gln Arg Arg Leu Gln
20 25
Phe Leu Ala Thr Ala Thr Phe Tyr Asp Ile Ala Met Thr
30 35
Ala Gly Tyr Leu Val Ala Ala Val Arg Phe Tyr 45
50 55
Glu Lys Gly Thr His Arg Gly Leu Tyr Ser Ile Gln Lys Thr
60 65
Leu Lys Phe Phe Gln Val Leu Gln Ile Val His 75
80 85
Leu Ile Gly Ile Val Pro Thr Ser Val Ile Val Thr Gly Val Gln
90 95
Val Ser Ser Arg Val Phe Met Val Trp Ile Thr His Ser 105
110 115
Lys Pro Ile Gln Arg Gln Ser Val Val Leu Phe Leu Val Ala
120 125
Trp Thr Val Thr Thr Arg Tyr Val Phe Thr Phe 135
140 145
Leu Leu Asp His Leu Pro Tyr Phe Ile Lys Trp Ala Arg Asn
150 155
Phe Phe Ile Leu Leu Tyr Pro Val Gly Val Ala Gly Gln Leu
160 165
Phe Thr Val Thr Thr Thr Thr Thr Thr Thr Thr Thr Thr Thr
170 175
Ser Ile Arg Leu Pro Asn Lys Tyr Asn Val Ser Phe Asp Tyr Tyr
185 190
Tyr Phe Leu Leu Ile Thr Met Ala Ser Tyr Ile Pro Leu Phe Pro
195 200
Gln Leu Tyr Phe His Met Leu Arg Gln Arg Arg Lys Val Leu His
205 210
Gly Gln Val Ile Val Lys Asp Asp 215 220 225 230 235 240

<210> 9
<211> 166
<212> PRT
<213> Homo sapiens
<220>
<221> misc_Feature
<222> Inocyte ID No: 151133C01
<400> 9
Met Ala Asp Asp Asp Val Leu Phe Gln Asp Val Tyr Gln Leu Cys
1 5
Glu Val Ile Gly 10 15
15 20 25 30 35 40 45 50 55 60 65 70 75 80 85 90 95 100 105 110 115 120 125 130 135 140 145 150 155 160 165 170 175 180 185 190 195 200 205 210 215 220 225 230 235 240

9/24

WO 0065333

PCT/US00/0777

PF-0683 PCT

<210> 10
<211> 354
<212> PRT
<213> Homo sapiens
<220>
<221> misc_Feature
<222> Inocyte ID No: 167761C01
<400> 10
Met Asn Ile Ala Arg Arg Lys Gln Gln Met Lys Asp Met Ile
1 5
Val Gln Thr Leu Asn Thr Met Lys Gln Gln Leu Leu Asp Asp Ala
10 15
Thr Asn Met Gln Lys Asp Val Ile 20 25
Pro Val Gly Thr Arg Gln Ile Lys Cys Cys Ile Arg Gln Ile Gln
30 35
Glu Leu Ile Ile Ser Arg Leu Asn Gln 40 45
50 55
Glu Leu Ile Ile Ser Arg Leu Asn Gln 60 65
70 75
Ile Ser Ser Val Asp Tyr Leu Arg Gln Ser Phe Val Gly Thr Leu
80 85
Glu Arg Cys Leu Asp Ser Leu Gln Lys Gln Asp Val Ser Val
90 95
His Ile Thr Ser Asn Tyr Leu Lys Gln Ile Leu Asn Ala Tyr
100 105
His Val Gln Val Phe His Ser Gly Ser Ser Val Thr Arg Met
110 115
Leu Trp Gln Gln Ile Lys Gln Ile Ile Gln Arg Ile Thr Trp Val
120 125
Ser Pro Pro Ala Ile Thr Leu Gln Trp Lys Arg Lys Val Ala
130 135
Glu Ala Ile Gln Ser Leu Ser Ala Ser Leu Asn Lys Ser 140 145
150 155
Cys Ser Gln Phe Arg Thr Arg Leu Asn Ser His Gln Ala Phe
160 165
Ala Ala Ser Leu Arg Gln Leu Gln Ala Gly His Ser Gly Arg 170 175
180 185
Glu Lys Thr Gln Asp Leu Trp Leu Arg Val Arg Lys Asp His Ala
190 195
Pro Arg Leu Ala Arg Leu Ser Leu Gln Ser Cys Ser Leu Gln 200 205
210 215
Val Leu Leu His Arg Lys Pro Lys Leu Gln Gln Gln Gly Arg 220 225 230 235 240

10/24

PF-0683 PCT

PF-0683 PCT

245
Gly Glu Tyr Gly Val Val Tyr Leu Cys Asn Trp Gly Gly His
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520

His Met Ala Pro Glu Leu Phe Thr Gly Lys Tyr
Asp Asn Ser Val Asp Val Tyr Ala Phe Gly Ile Leu Thr Trp Tyr
Ile Cys Ser Gly Ser Val Lys Leu Pro Glu Ala Phe Glu Arg
Ala Ser Lys Asp His Leu Trp Asn Val Arg Arg Gly Ala Arg
Pro Glu Arg Leu Val Phe Asp Glu Cys Trp Glu Leu
Glu Ala Cys Trp Asp Gly Asp Pro Leu Lys Asp Pro Leu Gly
Ile Val Glu Pro Leu Glu Ile Lys Asn Arg Leu Cys
Ser Asn Ser Glu Glu Pro Asn Arg Gly Leu Asp Asp Ser Thr

<210> 11

<211> 58

<212> PRT

<213> Homo sapiens

<214>

<215> misc. feature

<216> INCYTE ID No: 1270442CD1

<400> 11

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Met Arg Leu Arg Glu Arg Ser Leu Arg Glu Asp Pro Asp Leu Arg
Glu Glu Leu Ala Ser Leu Ala Arg Gly Cys Asp Phe Val Leu Pro
Ser Arg Phe Lys Arg Leu Lys Ala Phe Glu Val Glu Thr
Arg Lys Glu Glu Pro Leu Pro Ala Thr Ser Glu Ser Ile Pro
Thr Thr Phe Pro Glu Glu Glu Glu Glu Glu Glu Glu Glu Glu
Asp Ala Val Ile Ser Lys Ile Glu Ser Thr Phe Ala Arg Phe Pro
Pro

11/24

12/24

<210> 12

<211> 142

<212> PRT

PF-0683 PCT

<213> Homo sapiens

<220> misc_feature

<221> Incode ID No: 1877133CD1

<400> 12

Met Ser Thr Ala Arg Val Pro Ala Asp Lys Pro Val Arg Ile

Ala Phe Ser Leu Asn Asp Ala Ser Asp Asp Thr Pro Gln Asp

Ser Ile Pro Leu Val Phe Pro Gln Leu Asp Gln Leu Gln Pro

Leu Pro Pro Cys His Asp Ser Gln Asp Ser Met Gln Val Phe Lys

Gln His Cys Gln Ile Ala Gln Gln Tyr Thr Gln Val Lys Lys Gln

Ile Thr Leu Leu Gln Gln Arg Lys Lys Val Leu Ile Ala Lys

Asp Gln Ala Gln Lys Gln Lys Val Asp Ala Ala Gln Leu Val

Gln Phe Gln Ala Leu Thr Gln Gln Asn Arg Thr Leu Arg Leu Ala

Gln Ser Gln Cys Val Gln Gln Leu Gln Lys Leu Arg Ile Gln Tyr

Gln Lys Arg Gln Gln Ser Ser

140

140

140

140

140

140

140

140

140

140

140

140

140

140

140

140

140

140

140

WO 005332
PF-0683 PCT

<213> Homo sapiens

<220> misc_feature

<221> Incode ID No: 216815CD1

<400> 14

Met Ser Ile Ala Gln Val Ala Ala Gln Gln Ile Arg Val Pro Leu

Lys Thr Gln Phe Leu His Asn Gln Arg Ala Met Gln Asn Met Arg

Lys Thr Tyr Trp Ser Ser Arg Ser Gln Phe Lys Asn Asn Phe Leu

Asn Ile Asp Pro Ile Thr Met Ala Tyr Ser Leu Asn Ser Ser 45

Gln Gln Arg Leu Ile Pro Leu Gln His Ala Ser Lys Ser Ala Pro

Met Asn Gln His Cys Phe Ala Gln Asn Gln Pro Ser Gln Lys Ser

Ser Leu Pro Leu Leu Ile Pro Ser Gln Asn Leu Gln Pro

His Gln Gln Asp Gln Val Val Cys Gln Phe Lys Lys Leu Thr Val

Asn Gln Val Cys Ala Ser Thr Pro Leu Thr Pro Ile Lys Asn

Ser Pro Ser Leu Phe Pro Cys Ala Ser Thr Pro Leu Cys Gln Arg Gln

Arg Pro Leu Pro 155 160 165 170 175 180 185 190 195 200

195

195

195

195

195

195

195

195

195

195

195

195

195

195

195

195

195

195

WO 005332

PF-0683 PCT

<213> Homo sapiens

<220> misc_feature

<221> Incode ID No: 216815CD1

<400> 14

Met Ser Ile Ala Gln Val Ala Ala Gln Gln Ile Arg Val Pro Leu

Lys Thr Gln Phe Leu His Asn Gln Arg Ala Met Gln Asn Met Arg

Lys Thr Tyr Trp Ser Ser Arg Ser Gln Phe Lys Asn Asn Phe Leu

Asn Ile Asp Pro Ile Thr Met Ala Tyr Ser Leu Asn Ser Ser 45

Gln Gln Arg Leu Ile Pro Leu Gln His Ala Ser Lys Ser Ala Pro

Met Asn Gln His Cys Phe Ala Gln Asn Gln Pro Ser Gln Lys Ser

Ser Leu Pro Leu Leu Ile Pro Ser Gln Asn Leu Gln Pro

His Gln Gln Asp Gln Val Val Cys Gln Phe Lys Lys Leu Thr Val

Asn Gln Val Cys Ala Ser Thr Pro Leu Thr Pro Ile Lys Asn

Ser Pro Ser Leu Phe Pro Cys Ala Ser Thr Pro Leu Cys Gln Arg Gln

Arg Pro Leu Pro 155 160 165 170 175 180 185 190 195 200

195

195

195

195

195

195

195

195

195

195

195

195

195

195

195

195

195

195

PF-0683 PCT

<400> 23
 agctgcctca ggtatcagaa aagagctct tcaatgtct aagagagat atccacaaag 60
 aactctgcca aactattct gaaatattc tgaatgtct caagatcca tcaatctaa 120
 gttacatgaa aagatgaaa agatgaaa cttatgaaa cttatgaaa cttatgaaa 180
 gttacatgaa aagatgaaa agatgaaa cttatgaaa cttatgaaa cttatgaaa 240
 tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa 300
 tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa 360
 tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa 420
 tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa 480
 tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa 540
 tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa tttacatgaa 594

<210> 24
 <211> 2480
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc feature
 <222> Inqze ID No: 167743C01

<400> 24
 ccaaaatc tgaatcag tcaaaagaa gaaatcagat gtaataatc atgaatgat 60
 atctgcacac gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 120
 aagagagac caatgagac caatgagac aatgcagca tccagatgat tccagatgat 180
 atactcttc gactatcaa gaaatcagat aatgcagca tccagatgat gaaatcagat 240
 aagatgaaa aagatgaaa aagatgaaa aagatgaaa aagatgaaa aagatgaaa 300
 gaaatcagat tttacatgaa gttccatgt aagatgaaa tcaatgaaa aatcaaaag 360
 atactcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 420
 gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 480
 gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 540
 gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 594

<210> 25
 <211> 1679
 <212> Homo sapiens

<220>
 <221> misc feature
 <222> Inqze ID No: 1270442C01

<400> 25
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 60
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 120
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 180
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 240
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 300
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 360
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 420
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 480
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 540
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 594

PF-0683 PCT

<400> 25
 <210> 25
 <211> 1679
 <212> Homo sapiens

<220>
 <221> misc feature
 <222> Inqze ID No: 1270442C01

<400> 25
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 60
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 120
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 180
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 240
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 300
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 360
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 420
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 480
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 540
 gttatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 594

<210> 26
 <211> 1053
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc feature
 <222> Inqze ID No: 167743C01

<400> 26
 aagatgaaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 60
 aagatgaaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 120
 aagatgaaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 180
 aagatgaaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 240
 aagatgaaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 300
 aagatgaaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 360
 aagatgaaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 420
 aagatgaaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 480
 aagatgaaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 540
 aagatgaaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa gaaatcagaa 594

INTERNATIONAL SEARCH REPORT

<p>INTERNATIONAL APPLICATION No. PCT/US 00/07277</p>	
<p>C (Classification documents considered to be relevant)</p>	<p>Relevant to claim No.</p>
<p>P. X TANQUE, T. ET AL.: "Molecular Cloning and Characterization of a Novel Dual Specificity Phosphatase Kinase (p38) FROM <i>PHOSPHOLIPASE C</i> vol. 274, no. 28, 9 July 1999 (1999-07-09), pages 1949-1956, XP002146678 the whole document</p>	<p>1-3, 5, 6, 8-14</p>
<p>A GROHN, L.A. ET AL.: "Differential regulation of the Mnk, Skp and Rk/p38 kinases by Pyx1, a novel cytosolic dual-specific phosphatase" vol. 15, no. 14, 15 July 1996 (1996-07-15), pages 3621-3632, XP00925967 the whole document</p>	<p>1-17, 20, 23</p>
<p>A MO 99 01541 A (TULARIK INC.) 14 January 1999 (1999-01-14) Abstract, line 10 -page 16, line 13 -page 17 -page 18; claims 1-11</p>	<p>1-17, 20, 23</p>
<p>A MO 99 00507 A (INCYTE PHARMACEUTICALS, INC.) 17 January 1999 (1999-01-07) page 2, line 13 -page 3, line 31 page 38, line 10 -page 46, line 10 page 55 -page 57; claims 1-21</p>	<p>1-17, 20, 23</p>

2

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 00/07277

<p>Box 1 Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)</p>	
<p>This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:</p>	
<p>1. <input checked="" type="checkbox"/> Claims Nos. 18, 19, 21 and 22 because they relate to subject matter not required to be searched by the Authority, namely:</p>	
<p>Although claim 16 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.</p>	
<p>2. <input checked="" type="checkbox"/> Claims Nos. 18, 19, 21 and 22 because they relate to some of the International Application may not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out. See FURTHER INFORMATION SHEET PCT/ISA/210</p>	
<p>3. <input type="checkbox"/> Claims Nos. because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 8.4(a).</p>	
<p>Box 2 Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)</p>	
<p>This International Searching Authority found multiple inventions in this International Application, as follows:</p>	
<p>1. <input type="checkbox"/> As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.</p>	
<p>2. <input type="checkbox"/> As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.</p>	
<p>3. <input type="checkbox"/> As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos. 1-17, 20, 23 partially</p>	
<p>4. <input checked="" type="checkbox"/> No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention set mentioned in the claims; it is covered by claim Nos. 1-17, 20, 23 partially</p>	
<p>Remise en Protest <input type="checkbox"/> The additional search fees were accompanied by the applicant's protest. <input type="checkbox"/> No protest accompanied the payment of additional search fees.</p>	

1. Claims: 1-17, 20, 23 partially

Invention 1

An isolated polypeptide comprising an amino acid sequence selected from the group consisting of: a) an amino acid sequence having the SEQ ID NO: 1, b) a naturally occurring amino acid sequence having at least 98% sequence identity to SEQ ID NO: 1, c) a biologically active fragment of SEQ ID NO: 1, d) an immunogenic fragment of SEQ ID NO: 1, an isolated polynucleotide encoding a polypeptide having an amino acid sequence having the amino acid sequence of a polypeptide, a cDNA transformed with said recombinant polynucleotide; a transgenic organism comprising said recombinant polynucleotide; a method for producing said polypeptide; an isolated antibody which specifically binds to said polypeptide; an isolated polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of: a) a polynucleotide sequence having the SEQ ID NO: 15, b) a naturally occurring polynucleotide sequence having at least 98% sequence identity to the SEQ ID NO: 15, c) a biologically active fragment of the SEQ ID NO: 15, d) an immunogenic fragment of the SEQ ID NO: 15, e) a polynucleotide sequence complementary to a) - d), an isolated polynucleotide sequence complementary to a) - d), an equivalent of a) - d); a method for detecting a target polynucleotide in a sample having the sequence of said polynucleotide by hybridizing with a probe; a pharmaceutical composition comprising an effective amount of said polypeptide; a method for treating a disease or condition associated with decreased expression of functional Hsp70, comprising administering to a patient said pharmaceutical composition; a method for screening a compound for effectiveness as an agonist or antagonist of said polypeptide; a method for screening a compound for effectiveness in altering expression of a polynucleotide having the SEQ ID NO: 15;

2. Claims: 1-17, 20, 23 partially

Invention 2

Idem as subject 1 but limited to SEQ ID NOS: 2 and 16;

3.-14. Claims: 1-17, 20, 23 partially

Inventions 3-14

Idem as subject 1 but limited to SEQ ID NOS: 3-14 and 17-28

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 216

Continuation of Box 1.2

Claims Nos.: 18, 19, 21 and 22

No search can be carried out for such purely speculative claims whose wording is, in fact, a mere recitation of the result to be achieved.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO Board when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following the receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.
PCT/US 00/07277

Patent (doc. no.) cited in search report	Publication date	Patent family members)	Publication date
WO 9839446 A	11-09-1998	AU 6546388 A EP 0972029 A EP 0972030 A WO 9839448 A	22-09-1998 19-01-2000 19-01-2000 11-09-1998
WO 9901541 A	14-01-1999	AU 726294 B AU 828459 A EP 1006693 A	02-11-2000 25-01-1999 07-06-2000
WO 9900507 A	07-01-1999	US 5955338 A AU 8269998 A EP 0996733 A	21-09-1999 19-01-1999 03-05-2000